

Rabbit CBL Polyclonal Antibody | anti-CBL antibody

CBL Polyclonal Antibody

CATALOG # AAA280350	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

CBL; Polyclonal Antibody; CBL Polyclonal Antibody; anti-CBL antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	906
Applicable Applications for anti-CBL antibody	IHC (Immunohistochemistry), WB (Western Blot)
Route	Synthetic Peptide
Immunogen	A synthetic peptide of human CBL
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


IMAGES


IHC (Immunohistochemistry)


Immunohistochemistry of paraffin-embedded mouse liver using CBL antibody (40x lens).


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat kidney using CBL antibody (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse stomach using CBL antibody (40x lens).



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using CBL antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-CBL antibody

This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia. Mutations in this gene are also the cause of Noonan syndrome-like disorder.

NCBI AND UNIPROT INFORMATION

NCBI GI #	52426745
NCBI GeneID	867
NCBI Accession #	NP_005179.2
NCBI GenBank Nucleotide #	NM_005188.3
UniProt Accession #	P22681; A3KMP8
Molecular Weight	100kDa
NCBI Official Full Name	E3 ubiquitin-protein ligase CBL
NCBI Official Synonym Full Names	Cbl proto-oncogene, E3 ubiquitin protein ligase
NCBI Official Symbol	CBL
NCBI Official Synonym Symbols	CBL2; NSLL; C-CBL; RNF55; FRA11B
NCBI Protein Information	E3 ubiquitin-protein ligase CBL
UniProt Protein Name	E3 ubiquitin-protein ligase CBL
UniProt Gene Name	CBL
UniProt Synonym Gene Names	CBL2; RNF55
UniProt Entry Name	CBL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.