

Rabbit BAG2 Polyclonal Antibody | anti-BAG2 antibody

BAG2 Polyclonal Antibody

CATALOG # AAA280404	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

BAG2; Polyclonal Antibody; BAG2 Polyclonal Antibody; anti-BAG2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	211
Applicable Applications for anti-BAG2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human BAG2
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney using BAG2 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using BAG2 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 60s.

ADDITIONAL INFORMATION

Related Product Information for anti-BAG2 antibody	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757834
NCBI GeneID	9532
NCBI Accession #	NP_004273.1
NCBI GenBank Nucleotide #	NM_004282.3
UniProt Accession #	O95816; Q08AS9; Q6FID0; B4DXE2
Molecular Weight	24kDa
NCBI Official Full Name	BAG family molecular chaperone regulator 2
NCBI Official Synonym Full Names	BCL2-associated athanogene 2
NCBI Official Symbol	BAG2
NCBI Official Synonym Symbols	BAG-2; dJ41711.2
NCBI Protein Information	BAG family molecular chaperone regulator 2
UniProt Protein Name	BAG family molecular chaperone regulator 2
UniProt Gene Name	BAG2
UniProt Synonym Gene Names	BAG-2
UniProt Entry Name	BAG2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bag2-antibody-immunohistochemistry-western-blot-elisa/280404>