

Rabbit anti-Human ATP7A Polyclonal Antibody | anti-ATP7A antibody

ATP7A Polyclonal Antibody

CATALOG # AAA280571	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot
-------------------------------	-----------------------	----------------------------	-------------------------------------


PRODUCT SYNONYMS

ATP7A; Polyclonal Antibody; ATP7A Polyclonal Antibody; anti-ATP7A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	1500
Applicable Applications for anti-ATP7A antibody	WB (Western Blot)
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human ATP7A
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using ATP7A antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver injury using ATP7A antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat kidney using ATP7A antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of 293T cells, using ATP7A antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-ATP7A antibody

This gene encodes a transmembrane protein that functions in copper transport across membranes. This protein is localized to the trans Golgi network, where it is predicted to supply copper to copper-dependent enzymes in the secretory pathway. It relocates to the plasma membrane under conditions of elevated extracellular copper, and functions in the efflux of copper from cells. Mutations in this gene are associated with Menkes disease, X-linked distal spinal muscular atrophy, and occipital horn syndrome. Alternatively-spliced transcript variants have been observed.

NCBI AND UNIPROT INFORMATION

NCBI GI #	532691750
NCBI GeneID	538
NCBI Accession #	NP_000043.4
NCBI GenBank Nucleotide #	NM_000052.6
UniProt Accession #	Q04656; O00227; O00745; Q9BY8; B1AT72
Molecular Weight	163kDa
NCBI Official Full Name	copper-transporting ATPase 1 isoform 1
UniProt Protein Name	Copper-transporting ATPase 1
UniProt Gene Name	ATP7A
UniProt Synonym Gene Names	MC1; MNK
UniProt Entry Name	ATP7A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/atp7a-antibody-western-blot-elisa/280571>