

Rabbit anti-Human ABCC1 Polyclonal Antibody | anti-ABCC1 antibody

ABCC1 Polyclonal Antibody

CATALOG # AAA280795	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot
-------------------------------	-----------------------	----------------------------	-------------------------------------


PRODUCT SYNONYMS

ABCC1; Polyclonal Antibody; ABCC1 Polyclonal Antibody; ABC29; ABCC; GS-X; MRP; MRP1; anti-ABCC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	1531
Applicable Applications for anti-ABCC1 antibody	WB (Western Blot)
Immunogen	Recombinant protein of human ABCC1
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES





IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human gastric cancer using ABCC1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using ABCC1 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver using ABCC1 antibody at dilution of 1:100 (40x lens).

IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat heart using ABCC1 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of HepG2 cells, using ABCC1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.

ADDITIONAL INFORMATION

Related Product Information for anti-ABCC1 antibody	The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra-and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This full transporter is a member of the MRP subfamily which is involved in multi-drug resistance. This protein functions as a multispecific organic anion transporter, with oxidized glutathione, cysteinyl leukotrienes, and activated aflatoxin B1 as substrates. This protein also transports glucuronides and sulfate conjugates of steroid hormones and bile salts. Alternatively spliced variants of this gene have been described but their full-length nature is unknown.
Product Categories/Family for anti-ABCC1 antibody	Primary antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	296439301
NCBI GeneID	4363
UniProt Accession #	P33527 ; O14819 ; O43333 ; P78419 ; Q59GI9 ; Q9UQ97 ; Q9UQ99 ; Q9UQA0 ; A3RJX2 ; C9JPJ4
Molecular Weight	172kDa
NCBI Official Full Name	Multidrug resistance-associated protein 1
NCBI Official Synonym Full Names	ATP binding cassette subfamily C member 1

NCBI Official Symbol	ABCC1
NCBI Official Synonym Symbols	MRP; ABCC; GS-X; MRP1; ABC29
NCBI Protein Information	multidrug resistance-associated protein 1
UniProt Protein Name	Multidrug resistance-associated protein 1
UniProt Gene Name	ABCC1
UniProt Synonym Gene Names	MRP; MRP1; LTC4 transporter

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/abcc1-antibody-western-blot-elisa/280795>