

Rabbit CDK1 Polyclonal Antibody | anti-CDK1 antibody

CDK1 Polyclonal Antibody

CATALOG # AAA280870	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

CDK1; Polyclonal Antibody; CDK1 Polyclonal Antibody; CDC2; CDC28A; P34CDC2; anti-CDK1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	297
Applicable Applications for anti-CDK1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human CDK1
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse brain using CDK1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human esophagus using CDK1 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human lung cancer using CDK1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat brain using CDK1 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CDK1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-CDK1 antibody

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Product Categories/Family for anti-CDK1 antibody

[Primary antibody](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	334302921
NCBI GeneID	983
UniProt Accession #	P06493 ; O60764 ; A8K7C4 ; C9J497
Molecular Weight	34kDa
NCBI Official Full Name	Cyclin-dependent kinase 1
NCBI Official Synonym Full Names	cyclin dependent kinase 1

NCBI Official Symbol	CDK1
NCBI Official Synonym Symbols	CDC2; CDC28A; P34CDC2
NCBI Protein Information	cyclin-dependent kinase 1
UniProt Protein Name	Cyclin-dependent kinase 1
UniProt Gene Name	CDK1
UniProt Synonym Gene Names	CDC2; CDC28A; CDKN1; P34CDC2; CDK1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cdk1-antibody-immunohistochemistry-western-blot-elisa/280870>