

Rabbit anti-Human, Mouse TIMP1 Polyclonal Antibody | anti-TIMP1 antibody

TIMP1 Polyclonal Antibody

CATALOG # AAA280954	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Western Blot
-------------------------------	-----------------------	-----------------------------------	-------------------------------------


PRODUCT SYNONYMS

TIMP1; Polyclonal Antibody; TIMP1 Polyclonal Antibody; CLGI; EPA; EPO; HCI; TIMP; TIMP-1; anti-TIMP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence Length	207
Applicable Applications for anti-TIMP1 antibody	WB (Western Blot)
Immunogen	Recombinant protein of human TIMP1
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of extracts of mouse lung, using TIMP1 antibody at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using TIMP1 antibody at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-TIMP1 antibody	This gene belongs to the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction.
Product Categories/Family for anti-TIMP1 antibody	Primary antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	135850
NCBI GeneID	7076
UniProt Accession #	P01033; Q14252; Q9UCU1
Molecular Weight	23kDa
NCBI Official Full Name	Metalloproteinase inhibitor 1
NCBI Official Synonym Full Names	TIMP metallopeptidase inhibitor 1
NCBI Official Symbol	TIMP1
NCBI Official Synonym Symbols	EPA; EPO; HCI; CLGI; TIMP; TIMP-1
NCBI Protein Information	metalloproteinase inhibitor 1
UniProt Protein Name	Metalloproteinase inhibitor 1
UniProt Gene Name	TIMP1
UniProt Synonym Gene Names	CLGI; TIMP; EPA; Collagenase inhibitor; TIMP-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/timp1-antibody-western-blot-elisa/280954>