

Rabbit anti-Mouse TPM3 Polyclonal Antibody | anti-TPM3 antibody

TPM3 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA281103	Rabbit	Mouse	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TPM3; Polyclonal Antibody; TPM3 Polyclonal Antibody; CAPM1; CFTD; HEL-189; HEL-S-82p; hscp30; NEM1; OK/SW-cl.5; TM-5; TM3; TM30; TM30nm; TM5; TPMsk3; TRK; anti-TPM3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Sequence	MMEAIKKKMQ MLKLDKENAL DRAEQAEAEQ KQAEERSKQL EDELAAMQKK LKGTEDELDK YSEALKDAQE KLELAEEKKAA DAAEAVASLN RRIQLVEEEL DRAQERLATA LQKLEEAeka ADESERGMKV IENRALKDEE KMELQEIQLK EAKHIAEEAD RKYEEVARKL VIIEGDLERT EERAELAESK CSELEEELKN VTNNLKSLEA QAEKYSQKED KYEEEEIKILT DKLKEAETRA EFAERSVAKL EKTID
Sequence Length	248
Applicable Applications for anti-TPM3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human TPM3
Immunogen Species	Human
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Cellular Location	Cytoplasm, cytoskeleton
Positive Samples	Mouse skeletal muscle, Mouse lung, Mouse heart, Mouse spinal cord
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded mouse heart using TPM3 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using TPM3 antibody at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.Lysates/proteins: 25ug per lane.Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-TPM3 antibody	This gene encodes a member of the tropomyosin family of actin-binding proteins. Tropomyosins are dimers of coiled-coil proteins that provide stability to actin filaments and regulate access of other actin-binding proteins. Mutations in this gene result in autosomal dominant nemaline myopathy and other muscle disorders. This locus is involved in translocations with other loci, including anaplastic lymphoma receptor tyrosine kinase (ALK) and neurotrophic tyrosine kinase receptor type 1 (NTRK1), which result in the formation of fusion proteins that act as oncogenes. There are numerous pseudogenes for this gene on different chromosomes. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-TPM3 antibody	Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	114155144
NCBI GeneID	7170
NCBI Accession #	NP_001036816.1
NCBI GenBank Nucleotide #	NM_001043351.1
UniProt Accession #	P06753; P12324; Q2QD06; Q5VU58; Q5VU63; Q5VU66; Q5VU71; Q5VU72; Q8TCG3; Q969Q2; Q9NQH8; D3DV71
Molecular Weight	Calculated: 18kDa; 28kDa; 29kDa; 32kDa Observed: 37kDa
NCBI Official Full Name	tropomyosin alpha-3 chain isoform Tpm3.2cy
NCBI Official Synonym Full Names	tropomyosin 3
NCBI Official Symbol	TPM3
NCBI Official Synonym Symbols	TM3; TM5; TRK; CFTD; NEM1; TM-5; TM30; CAPM1; TM30nm; TPMsk3; hscp30; HEL-189; HEL-S-82p; OK/SW-cl.5
NCBI Protein Information	tropomyosin alpha-3 chain
UniProt Protein Name	Tropomyosin alpha-3 chain
UniProt Gene Name	TPM3
UniProt Synonym Gene Names	hTM5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tpm3-antibody-immunohistochemistry-western-blot-elisa/281103>