

Rabbit TBP Polyclonal Antibody | anti-TBP antibody

TBP Polyclonal Antibody

CATALOG # AAA281470	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

TBP; Polyclonal Antibody; TBP Polyclonal Antibody; GTF2D; GTF2D1; HDL4; SCA17; TFIID; TATA-box-binding protein; anti-TBP antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Concentration	1.69 mg/ml (varies by lot)
Sequence Length	319
Applicable Applications for anti-TBP antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 160-319 of human TBP (NP_001165556.1).
Immunogen Sequence	LKTIALRRARNAEYNPKRFAAVIMRIREPRTTALIFSSGKMVCTGAKSEEQSRLAARKYARVVQKLGFPKFLDFKIQNMVGS CDVKFPIRLEGLVLTHQQFS
Cellular Location	Nucleus
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES



IF (Immunofluorescence)
Immunofluorescence-TBP Polyclonal Antibody



IF (Immunofluorescence)
Immunofluorescence-TBP Polyclonal Antibody



IF (Immunofluorescence)
Immunofluorescence-TBP Polyclonal Antibody

ADDITIONAL INFORMATION

Related Product Information for anti-TBP antibody	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes TBP, the TATA-binding protein. A distinctive feature of TBP is a long string of glutamines in the N-terminus. This region of the protein modulates the DNA binding activity of the C terminus, and modulation of DNA binding affects the rate of transcription complex formation and initiation of transcription. The number of CAG repeats encoding the polyglutamine tract is usually 25-42, and expansion of the number of repeats to 45-66 increases the length of the polyglutamine string and is associated with spinocerebellar ataxia 17, a neurodegenerative disorder classified as a polyglutamine disease. Two transcript variants encoding different isoforms have been found for this gene.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	285026520
NCBI GeneID	6908
NCBI Accession #	NP_001165556.1
NCBI GenBank Nucleotide #	NP_001165556.1
UniProt Accession #	P20226
Molecular Weight	35kDa; 37kDa
NCBI Official Full Name	TATA-box-binding protein isoform 2
NCBI Official Synonym Full Names	TATA-box binding protein
NCBI Official Symbol	TBP
NCBI Official Synonym Symbols	HDL4; GTF2D; SCA17; TFIID; GTF2D1
NCBI Protein Information	TATA-box-binding protein
UniProt Protein Name	TATA-box-binding protein
UniProt Gene Name	TBP
UniProt Synonym Gene Names	GTF2D1; TF2D; TFIID
UniProt Entry Name	TBP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tbp-antibody-immunofluorescence-western-blot-elisa/281470>