

Rabbit ERK2 Antibody | anti-ERK2 antibody

[KO Validated] ERK2 Rabbit mAb

CATALOG # AAA281562	HOST Rabbit	APPLICATIONS Western Blot
-------------------------------	-----------------------	-------------------------------------


PRODUCT SYNONYMS

ERK2; [KO Validated] ERK2 Rabbit mAb; anti-ERK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Isotype	IgG
Purity/Purification	Affinity purification
Applicable Applications for anti-ERK2 antibody	WB (Western Blot)
Crossreactivity	Human, Mouse, Rat
Immunogen	A synthesized peptide derived from human ERK2.
Buffer	PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using ERK2 antibody (AAA281562) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.



WB (Western Blot)

Western blot analysis of extracts from normal (control) and ERK2 knockout (KO) HeLa cells, using ERK2 antibody (AAA281562) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

ADDITIONAL INFORMATION

Related Product Information for anti-ERK2 antibody	This gene encodes a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation, this kinase translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. One study also suggests that this protein acts as a transcriptional repressor independent of its kinase activity. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Two alternatively spliced transcript variants encoding the same protein, but differing in the UTRs, have been reported for this gene. [provided by RefSeq, Jan 2014]
--	---

NCBI AND UNIPROT INFORMATION

NCBI GeneID	5594
UniProt Accession #	P28482
Molecular Weight	41KDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-antibody/erk2/281562>