

Rabbit MAPK1/MAPK3 Polyclonal Antibody | anti-MAPK1/MAPK3 antibody

MAPK1/MAPK3 Rabbit pAb

CATALOG # AAA281771	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

MAPK1/MAPK3; Polyclonal Antibody; MAPK1/MAPK3 Rabbit pAb; anti-MAPK1/MAPK3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-MAPK1/MAPK3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human MAPK1/MAPK3.
Positive Samples	293T, NIH/3T3
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat lung using MAPK1/MAPK3 Rabbit pAb at dilution of 1:200 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse heart using MAPK1/MAPK3 Rabbit pAb at dilution of 1:200 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human urothelial carcinoma using MAPK1/MAPK3 Rabbit pAb at dilution of 1:200 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using MAPK1/MAPK3 Rabbit pAb at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK1/MAPK3 antibody

Background: MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation, this kinase translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets.

NCBI AND UNIPROT INFORMATION

NCBI GI #	27469582
NCBI GeneID	5562
UniProt Accession #	Q13131; P54646
Molecular Weight	64,009 Da
NCBI Official Full Name	PRKAA1 protein
NCBI Official Synonym Full Names	protein kinase, AMP-activated, alpha 1 catalytic subunit
NCBI Official Symbol	PRKAA1
NCBI Official Synonym Symbols	AMPK; AMPKa1
NCBI Protein Information	5'-AMP-activated protein kinase catalytic subunit alpha-1; ACACA kinase; AMPK alpha 1; HMGCR kinase; AMPK subunit alpha-1; tau-protein kinase PRKAA1; acetyl-CoA carboxylase kinase; AMP -activate kinase alpha 1 subunit; hydroxymethylglutaryl-CoA reductase
UniProt Protein Name	5'-AMP-activated protein kinase catalytic subunit alpha-1
UniProt Gene Name	PRKAA1
UniProt Synonym Gene Names	AMPK1; AMPK subunit alpha-1; ACACA kinase; HMGCR kinase
UniProt Entry Name	AAPK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mapk1-mapk3-antibody-immunohistochemistry-western-blot-elisa/281771>