

Rabbit ATM Polyclonal Antibody | anti-ATM antibody

ATM Rabbit pAb

CATALOG # AAA281919	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

ATM; Polyclonal Antibody; ATM Rabbit pAb; AT1; ATA; ATC; ATD; ATDC; ATE; TEL1; TELO1; anti-ATM antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-ATM antibody	WB (Western Blot)
Immunogen	Recombinant protein of human ATM.
Cellular Location	Cytoplasmic vesicle, Nucleus
Positive Samples	293T
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of extracts from normal (control) and ATM knockout (KO) 293T cells, using ATM antibody at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 180s.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using ATM antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3min.

ADDITIONAL INFORMATION

Related Product Information for anti-ATM antibody	Background: The protein encoded by this gene belongs to the PI3/PI4-kinase family. This protein is an important cell cycle checkpoint kinase that phosphorylates; thus, it functions as a regulator of a wide variety of downstream proteins, including tumor suppressor proteins p53 and BRCA1, checkpoint kinase CHK2, checkpoint proteins RAD17 and RAD9, and DNA repair protein NBS1. This protein and the closely related kinase ATR are thought to be master controllers of cell cycle checkpoint signaling pathways that are required for cell response to DNA damage and for genome stability. Mutations in this gene are associated with ataxia telangiectasia, an autosomal recessive disorder.
Product Categories/Family for anti-ATM antibody	Epigenetics & Nuclear Signaling; Epigenetic writers and erasers of core Histones; Phosphorylation; Research; Epigenetics & Nuclear Signaling; DNA Damage & Repair; Research; Protein phosphorylation; Research; Signal Transduction; Kinase; Serine/threonine k

NCBI AND UNIPROT INFORMATION

NCBI GI #	187951615
NCBI GeneID	472
UniProt Accession #	Q13315; O15429; Q12758; Q16551; Q93007; Q9NP02; Q9UCX7; B2RNX5
Molecular Weight	350,714 Da
NCBI Official Full Name	Ataxia telangiectasia mutated
NCBI Official Synonym Full Names	ataxia telangiectasia mutated
NCBI Official Symbol	ATM
NCBI Official Synonym Symbols	AT1; ATA; ATC; ATD; ATE; ATDC; TEL1; TELO1
NCBI Protein Information	serine-protein kinase ATM; AT mutated; A-T mutated; TEL1, telomere maintenance 1, homolog
UniProt Protein Name	Serine-protein kinase ATM
UniProt Gene Name	ATM
UniProt Synonym Gene Names	A-T mutated
UniProt Entry Name	ATM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/atm-antibody-western-blot-elisa/281919>