

Rabbit GSTA1 Polyclonal Antibody | anti-GSTA1 antibody

GSTA1 Rabbit pAb

CATALOG # AAA281922	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

GSTA1; Polyclonal Antibody; GSTA1 Rabbit pAb; GST-epsilon; GST2; GSTA1-1; GTH1; anti-GSTA1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-GSTA1 antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-222 of human GSTA1 (NP_665683.1).
Cellular Location	Cytoplasm
Positive Samples	HepG2, Mouse kidney, Mouse stomach, Rat liver, Rat kidney
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of U-2 OS cells using GSTA1 Rabbit pAb at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of L929 cells using GSTA1 Rabbit pAb at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



IF (Immunofluorescence)

Immunofluorescence analysis of C6 cells using GSTA1 Rabbit pAb at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using GSTA1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-GSTA1 antibody

Background: This gene encodes a member of a family of enzymes that function to add glutathione to target electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins, and products of oxidative stress. This action is an important step in detoxification of these compounds. This subfamily of enzymes has a particular role in protecting cells from reactive oxygen species and the products of peroxidation. Polymorphisms in this gene influence the ability of individuals to metabolize different drugs. This gene is located in a cluster of similar genes and pseudogenes on chromosome 6. Alternative splicing results in multiple transcript variants.

NCBI AND UNIPROT INFORMATION

NCBI GI #	22091454
NCBI GeneID	2938
NCBI Accession #	NP_665683.1
NCBI GenBank Nucleotide #	NM_145740.3
UniProt Accession #	P08263; Q14750; Q5GHF8; Q5SZC1
Molecular Weight	25,631 Da
NCBI Official Full Name	glutathione S-transferase A1
NCBI Official Synonym Full Names	glutathione S-transferase alpha 1
NCBI Official Symbol	GSTA1
NCBI Official Synonym Symbols	GST2; GTH1; GSTA1-1
NCBI Protein Information	glutathione S-transferase A1; GST-epsilon; GST HA subunit 1; GST, class alpha, 1; GST class-alpha member 1; glutathione S-transferase 2; glutathione S-aryltransferase A1; glutathione S-alkyltransferase A1; S-(hydroxyalkyl)glutathione lyase A1; glutathione
UniProt Protein Name	Glutathione S-transferase A1
UniProt Gene Name	GSTA1
UniProt Entry Name	GSTA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.