

Rabbit Pan Lactic acid-Lysine Polyclonal Antibody

Pan Lactic acid-Lysine Rabbit pAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA282001	Rabbit	Human, Mouse, Rat, Other (Wide Range)	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Pan Lactic acid-Lysine; Polyclonal Antibody; Pan Lactic acid-Lysine Rabbit pAb; anti-Pan Lactic acid-Lysine antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Other (Wide Range)
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Applicable Applications for anti-Pan Lactic acid-Lysine antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human Pan Lactic acid-Lysine.
Positive Samples	HeLa, NIH/3T3, C6
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of MCF7 cells using Pan Lactic acid-Lysine Rabbit pAb(AAA282001) at a dilution of 1:25 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




WB (Western Blot)

Western blot analysis of lysates from C6 cells, using Pan Lactic acid-Lysine Rabbit pAb (AAA282001) at 1:400 dilution. C6 cells were treated by Sodium lactate(100mM) for 24h. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 60s.



WB (Western Blot)

Western blot analysis of lysates from NIH/3T3 cells, using Pan Lactic acid-Lysine Rabbit pAb (AAA282001) at 1:400 dilution. NIH/3T3 cells were treated by Sodium lactate (100mM) for 24h. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 60s.



WB (Western Blot)

Western blot analysis of lysates from MCF7 cells, using Pan Lactic acid-Lysine Rabbit pAb (AAA282001) at 1:400 dilution. MCF7 cells were treated by Sodium lactate (100mM) for 24h. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 60s.

ADDITIONAL INFORMATION

Product Categories/Family for anti-Pan Lactic acid-Lysine antibody

Other modified Antibody

NCBI AND UNIPROT INFORMATION

Molecular Weight

Observed MW: 35-130KDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/pan-lactic-acid-lysine-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/282001>