

Rabbit SLC5A2 Polyclonal Antibody | anti-SLC5A2 antibody

SLC5A2 Rabbit pAb

CATALOG # AAA282062	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

SLC5A2; Polyclonal Antibody; SLC5A2 Rabbit pAb; SGLT2; anti-SLC5A2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-SLC5A2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human SLC5A2.
Positive Samples	293T, Jurkat, Mouse heart, Rat brain
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of human kidney using SLC5A2 Rabbit pAb at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of rat kidney using SLC5A2 Rabbit pAb at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse kidney using SLC5A2 Rabbit pAb at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using SLC5A2 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using SLC5A2 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.

ADDITIONAL INFORMATION

Related Product Information for anti-SLC5A2 antibody

Background: This gene encodes a member of the sodium glucose cotransporter family which are sodium-dependent glucose transport proteins. The encoded protein is the major cotransporter involved in glucose reabsorption in the kidney. Mutations in this gene are associated with renal glucosuria. Two transcript variants, one protein-coding and one not, have been found for this gene.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507033
NCBI GeneID	6524
NCBI Accession #	NP_003032.1
NCBI GenBank Nucleotide #	NM_003041.3
UniProt Accession #	P31639
Molecular Weight	72,897 Da
NCBI Official Full Name	sodium/glucose cotransporter 2
NCBI Official Synonym Full Names	solute carrier family 5 (sodium/glucose cotransporter), member 2

NCBI Official Symbol	SLC5A2
NCBI Official Synonym Symbols	SGLT2
NCBI Protein Information	sodium/glucose cotransporter 2; Na(+)/glucose cotransporter 2; solute carrier family 5 member 2; low affinity sodium-glucose cotransporter; solute carrier family 5 (sodium/glucose transporter), member 2
UniProt Protein Name	Sodium/glucose cotransporter 2
UniProt Gene Name	SLC5A2
UniProt Synonym Gene Names	SGLT2; Na(+)/glucose cotransporter 2
UniProt Entry Name	SC5A2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/slc5a2-antibody-immunohistochemistry-western-blot-elisa/282062>