

Rabbit anti-Human, Mouse Rad23B Antibody | anti-Rad23B antibody

Rad23B Rabbit pAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA282305	Rabbit	Human, Mouse	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Rad23B; Rad23B Rabbit pAb; RAD23B; HHR23B; HR23B; P58; UV excision repair protein RAD23 homolog B; anti-Rad23B antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Sequence	ELSRV
Applicable Applications for anti-Rad23B antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human Rad23B (NP_002865.1).
Cellular Location	Cytoplasm, Nucleus
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using Rad23B antibody at dilution of 1:100. Blue: DAPI for nuclear staining.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse testis using Rad23B Rabbit pAb at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon using Rad23B Rabbit pAb at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using Rad23B antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-Rad23B antibody

The protein encoded by this gene is one of two human homologs of *Saccharomyces cerevisiae* Rad23, a protein involved in the nucleotide excision repair (NER). This protein was found to be a component of the protein complex that specifically complements the NER defect of xeroderma pigmentosum group C (XP-c) cell extracts in vitro. This protein was also shown to interact with, and elevate the nucleotide excision activity of 3-methyladenine-DNA glycosylase (MPG), which suggested a role in DNA damage recognition in base excision repair. This protein contains an N-terminal ubiquitin-like domain, which was reported to interact with 26S proteasome, and thus this protein may be involved in the ubiquitin mediated proteolytic pathway in cells. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Product Categories/Family for anti-Rad23B antibody

[Epigenetics & Nuclear Signaling](#); [DNA Damage & Repair](#); [Polyclonal Antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GeneID

5887

Molecular Weight

Calculated MW: 35kDa/43kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-human-mouse-antibody/rad23b/282305>