

Mouse anti-Human IgG (Fc) Polyclonal Antibody | anti-IgG antibody

human IgG (Fc) Mouse mAb

CATALOG # AAA282311	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

IgG (Fc); Polyclonal Antibody; human IgG (Fc) Mouse mAb; COB1; YAP; YAP2; YAP65; YKI; YAP1; anti-IgG antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300, 50% glycerol, pH7.3.
Sequence	EEAEIPREVI ERLARSQIHS IRDLQRLLEI DSVGSEDSLD TSLRAHGVHA TKHVPEKRPL PIRRKRSIEE AVPAVCKTRT VIYEIPRSQV
Applicable Applications for anti-IgG antibody	IHC (Immunohistochemistry)
Immunogen	A synthesized peptide derived from human human IgG (Fc).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human tonsil using human IgG (Fc) antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon carcinoma using human IgG (Fc) antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

ADDITIONAL INFORMATION

Related Product Information for anti-IgG antibody	As a monomeric immunoglobulin that is predominately involved in the secondary antibody response and the only isotype that can pass through the human placenta, Immunoglobulin G (IgG) is synthesized and secreted by plasma B cells, and constitutes 75% of serum immunoglobulins in humans. IgG antibodies protect the body against the pathogens by agglutination and immobilization, complement activation, toxin neutralization, as well as the antibody-dependent cell-mediated cytotoxicity (ADCC). IgG tetramer contains two heavy chains (50 kDa) and two light chains (25 kDa) linked by disulfide bonds, that is the two identical halves form the Y-like shape. IgG is digested by pepsin proteolysis into Fab fragment (antigen-binding fragment) and Fc fragment ("crystallizable" fragment). IgG1 is most abundant in serum among the four IgG subclasses (IgG1, 2, 3 and 4) and binds to Fc receptors (FcgammaR) on phagocytic cells with high affinity.
Product Categories/Family for anti-IgG antibody	Immunology & Inflammation; Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	77695917
NCBI GeneID	5154
NCBI Accession #	NP_002598.4
NCBI GenBank Nucleotide #	NM_002607.5
UniProt Accession #	P04085; B5BU73
Molecular Weight	24,043 Da
NCBI Official Full Name	platelet-derived growth factor subunit A isoform 1 preproprotein
NCBI Official Synonym Full Names	platelet-derived growth factor alpha polypeptide
NCBI Official Symbol	PDGFA
NCBI Official Synonym Symbols	PDGF1; PDGF-A
NCBI Protein Information	platelet-derived growth factor subunit A; PDGF-1; PDGF A-chain; PDGF subunit A; platelet-derived growth factor A chain; platelet-derived growth factor alpha chain; platelet-derived growth factor alpha isoform 2 preproprotein
UniProt Protein Name	Platelet-derived growth factor subunit A
UniProt Gene Name	PDGFA
UniProt Synonym Gene Names	PDGF1; PDGF subunit A
UniProt Entry Name	PDGFA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/igg-fc-antibody-immunohistochemistry-elisa/282311>