

Rabbit anti-Human P4HB Monoclonal Antibody | anti-P4HB antibody

P4HB Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA282702	Rabbit	Human	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

P4HB; Monoclonal Antibody; P4HB Rabbit mAb; DSI; GIT; PDI; PHDB; PDIA1; PO4DB; PO4HB; PROHB; CLCRP1; ERBA2L; P4Hbeta; anti-P4HB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.
Sequence	KLGETYKDHE NIVIAKMDST ANEVEAVKVH SFPTLKFFPA SADRTVIDYN GERTLDGFKK FLESGGQDGA GDDDDLEDLE EAEPDMEED DDQKAVKDEL
Applicable Applications for anti-P4HB antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Cross Reactivity	Human, Mouse, Rat
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 409-508 of human P4HB (P07237).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded Human placenta using P4HB Rabbit mAb (AAA282702) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



WB (Western Blot)

Western blot analysis of various lysates using P4HB Rabbit mAb (AAA282702) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.



WB (Western Blot)

Western blot analysis of various lysates using P4HB Rabbit mAb (AAA282702) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s.

ADDITIONAL INFORMATION

Related Product Information for anti-P4HB antibody	This gene encodes the beta subunit of prolyl 4-hydroxylase, a highly abundant multifunctional enzyme that belongs to the protein disulfide isomerase family. When present as a tetramer consisting of two alpha and two beta subunits, this enzyme is involved in hydroxylation of prolyl residues in procollagen. This enzyme is also a disulfide isomerase containing two thioredoxin domains that catalyze the formation, breakage and rearrangement of disulfide bonds. Other known functions include its ability to act as a chaperone that inhibits aggregation of misfolded proteins in a concentration-dependent manner, its ability to bind thyroid hormone, its role in both the influx and efflux of S-nitrosothiol-bound nitric oxide, and its function as a subunit of the microsomal triglyceride transfer protein complex.
Product Categories/Family for anti-P4HB antibody	Epigenetics Nuclear Signaling; RNA Binding; Cancer; Signal Transduction; Cell Biology Developmental Biology; Cytoskeleton; Extracellular Matrix; Endocrine Metabolism; Lipid Metabolism; Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	20070125
NCBI GeneID	5034
NCBI Accession #	NP_000909.2
NCBI GenBank Nucleotide #	NM_000918.3
UniProt Accession #	P07237; P30037; P32079; Q15205; Q6LDE5; B2RDQ2
Molecular Weight	Calculated MW: 57kDa Observed MW: 57kDa
UniProt Protein Name	Protein disulfide-isomerase
UniProt Gene Name	P4HB
UniProt Synonym Gene Names	ERBA2L; PDI; PDIA1; PO4DB; PDI
UniProt Entry Name	PDIA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/p4hb-antibody-immunohistochemistry-western-blot-elisa/282702>