

Rabbit anti-Human WASP Monoclonal Antibody | anti-WAS antibody

WASP Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA283101	Rabbit	Human	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot

PRODUCT SYNONYMS

WASP; Monoclonal Antibody; WASP Rabbit mAb; THC; IMD2; SCN_X; THC₁; WASPA; anti-WAS antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.
Sequence	MSGGPMGGRP GGRGAPAVQQ NIPSTLLQDH ENQRLFEMLG RKCLTLTAV VQLYLALPPG AEHWTKHECG AVCFVKDNPQ KSYFIRLYGL QAGRLLWEQE
Applicable Applications for anti-WAS antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Cross Reactivity	Human, Mouse, Rat
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human WASP (P42768).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of paraffin-embedded mouse spleen using WASP Rabbit mAb (AAA283101) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of paraffin-embedded rat spleen using WASP Rabbit mAb (AAA283101) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



IF (Immunofluorescence)

Immunofluorescence analysis of Jurkat cells using WASP Rabbit mAb (AAA283101) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of lysates from Mouse liver, using WASP Rabbit mAb (AAA283101) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



WB (Western Blot)

Western blot analysis of various lysates using WASP Rabbit mAb (AAA283101) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-WAS antibody	The Wiskott-Aldrich syndrome (WAS) family of proteins share similar domain structure, and are involved in transduction of signals from receptors on the cell surface to the actin cytoskeleton. The presence of a number of different motifs suggests that they are regulated by a number of different stimuli, and interact with multiple proteins. Recent studies have demonstrated that these proteins, directly or indirectly, associate with the small GTPase, Cdc42, known to regulate formation of actin filaments, and the cytoskeletal organizing complex, Arp2/3. Wiskott-Aldrich syndrome is a rare, inherited, X-linked, recessive disease characterized by immune dysregulation and microthrombocytopenia, and is caused by mutations in the WAS gene. The WAS gene product is a cytoplasmic protein, expressed exclusively in hematopoietic cells, which show signalling and cytoskeletal abnormalities in WAS patients. A transcript variant arising as a result of alternative promoter usage, and containing a different 5' UTR sequence, has been described, however, its full-length nature is not known.
Product Categories/Family for anti-WAS antibody	Signal Transduction; Cell Biology Developmental Biology; Cell Adhesion; Cytoskeleton; Actins; Immunology Inflammation; T Cell Receptor Signaling Pathway; Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507909
NCBI GeneID	7454
NCBI Accession #	NP_000368.1
NCBI GenBank Nucleotide #	NM_000377.2
UniProt Accession #	P42768; Q9BU11; Q9UNJ9
Molecular Weight	Calculated MW: 53kDa Observed MW: 60kDa
UniProt Protein Name	Wiskott-Aldrich syndrome protein
UniProt Gene Name	WAS
UniProt Synonym Gene Names	IMD2; WASp
UniProt Entry Name	WASP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/wasp-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/283101>