

Rabbit anti-Human RB-S780 Monoclonal Antibody | anti-RB1 antibody

Phospho-RB-S780 Rabbit mAb

CATALOG # AAA283112	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

RB-S780; Monoclonal Antibody; Phospho-RB-S780 Rabbit mAb; RB; pRb; OSRC; pp110; p105-Rb; PPP1R130; p110-RB1; Phospho-RB-S780; anti-RB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.
Sequence	TLSPIP
Applicable Applications for anti-RB1 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Cross Reactivity	Human, Mouse, Rat
Immunogen	A synthetic phosphorylated peptide around S780 of human RB (P06400).
Modification	Phospho S780
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded Mouse liver using Phospho-RB-S780 Rabbit mAb (AAA283112) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.




IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded Human colon using Phospho-RB-S780 Rabbit mAb (AAA283112) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded Rat pancreas using Phospho-RB-S780 Rabbit mAb (AAA283112) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



WB (Western Blot)

Western blot analysis of lysates from C6 cells, using Phospho-RB-S780 Rabbit mAb (AAA283112) at 1:1000 dilution. C6 cells were treated by 10% FBS at 37° for 30 minutes after serum-starvation overnight. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-RB1 antibody	The protein encoded by this gene is a negative regulator of the cell cycle and was the first tumor suppressor gene found. The encoded protein also stabilizes constitutive heterochromatin to maintain the overall chromatin structure. The active, hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene are a cause of childhood cancer retinoblastoma (RB), bladder cancer, and osteogenic sarcoma.
Product Categories/Family for anti-RB1 antibody	Epigenetics Nuclear Signaling; Transcription Factors; DNA Damage Repair; Protein phosphorylation; Cancer; Tumor suppressors; Cell Biology Developmental Biology; Cell Cycle; Cell cycle inhibitors; Cell Cycle Control-G1 S Checkpoint; Phosphorylated Antibodi

NCBI AND UNIPROT INFORMATION

NCBI GI #	108773787
NCBI GeneID	5925
NCBI Accession #	NP_000312.2
NCBI GenBank Nucleotide #	NM_000321.2
UniProt Accession #	P06400; P78499; Q5VW46; Q8IZL4; A8K5E3
Molecular Weight	Calculated MW: 106kDa Observed MW: 106kDa
UniProt Protein Name	Retinoblastoma-associated protein
UniProt Gene Name	RB1
UniProt Synonym Gene Names	Rb
UniProt Entry Name	RB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/rb-s780-antibody-immunohistochemistry-western-blot-elisa/283112>