

Rabbit anti-Human CYPOR Polyclonal Antibody | anti-POR antibody

CYPOR Rabbit pAb

CATALOG # AAA283174	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

CYPOR; Polyclonal Antibody; CYPOR Rabbit pAb; CPR; P450R; anti-POR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Sequence	MINMGDSHVD TSSTVSEAVA EEVSLFSMTD MILFSLIVGL LTYWFLFRKK KEEVPEFTKI QTLTSSVRES SFVEKMKKTG RNIIVFYGSQ TGTAEEFANR LSKDAHRYGM RGMSADPEEY DLADLSSLPE IDNALVVFCEM ATYGEDPTD NAQDFYDWLQ ETDVDLSGVK FAVFGLGNKT YEHFNAMGKY VDKRLEQLGA QRIFELGLGD DDGNLEEDFI TWREQFWPAV CEHFGVEATG EESSIRQYEL
Applicable Applications for anti-POR antibody	ELISA, WB (Western Blot)
Cross Reactivity	Human, Mouse, Rat
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human CYPOR (NP_000932.3).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of lysates from Mouse liver using CYPOR Rabbit pAb (AAA283174) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



WB (Western Blot)

Western blot analysis of lysates from HeLa cells using CYPOR Rabbit pAb (AAA283174) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s.

ADDITIONAL INFORMATION

Related Product Information for anti-POR antibody	This gene encodes an endoplasmic reticulum membrane oxidoreductase that is essential for multiple metabolic processes, including reactions catalyzed by cytochrome P450 proteins for metabolism of steroid hormones, drugs and xenobiotics. The encoded protein has a flavin adenine dinucleotide (FAD)-binding domain and a flavodoxin-like domain which bind two cofactors, FAD and FMN, that allow it to donate electrons directly from NADpHto all microsomal P450 enzymes. Mutations in this gene cause a complex set of disorders, including apparent combined P450C17 and P450C21 deficiency, amenorrhea and disordered steroidogenesis, congenital adrenal hyperplasia and Antley-Bixler syndrome, that resemble those caused by defects in steroid metabolizing enzymes such as aromatase, 21-hydroxylase, and 17 alpha-hydroxylase.
Product Categories/Family for anti-POR antibody	Cancer; Signal Transduction; Endocrine Metabolism; Mitochondrial metabolism; Lipid Metabolism; Lipases; Drug metabolism; Cardiovascular; Lipids; Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	127139033
NCBI GeneID	5447
NCBI Accession #	NP_000932.3
NCBI GenBank Nucleotide #	NM_000941.2
UniProt Accession #	P16435; Q16455; Q197M5; Q8N181; Q9H3M8; Q9UDT3
Molecular Weight	Calculated MW: 77kDa Observed MW: 77kDa
UniProt Protein Name	NADPH--cytochrome P450 reductase
UniProt Gene Name	POR
UniProt Synonym Gene Names	CPR; P450R
UniProt Entry Name	NCPR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cypor-antibody-western-blot-elisa/283174>