

Rabbit Rev-Erbalpha/NR1D1 Polyclonal Antibody | anti-NR1D1 antibody

Rev-Erbalpha/NR1D1 Rabbit pAb

CATALOG # AAA28372	HOST Rabbit	APPLICATIONS Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry, ELISA
------------------------------	-----------------------	---


PRODUCT SYNONYMS

Rev-Erbalpha/NR1D1; Polyclonal Antibody; Rev-Erbalpha/NR1D1 Rabbit pAb; NR1D1; EAR1; THRA1; THRAL; ear-1; hRev; nuclear receptor subfamily 1 group D member 1; anti-NR1D1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-NR1D1 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA
Application Notes	WB 1:500 - 1:2000 IHC-P 1:50 - 1:200 IF/ICC 1:50 - 1:200
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 280-360 of human Rev-Erb α /NR1D1 (NP_068370.1).
Species Reactivity/Cross Reactivity	Human, Mouse, Rat
Preparation and Storage	Store at -20°C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of PC-12 cells using Rev-Erb α /NR1D1 Rabbit pAb at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using Rev-Erb α /NR1D1 Rabbit pAb at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat liver using Rev-Erb α /NR1D1 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse brain using Rev-Erb α /NR1D1 antibody at dilution of 1:100 (40x lens).

**WB (Western Blot)**

Western blot analysis of lysates from Rat lung, using Rev-Erbα/NR1D1 Rabbit pAb at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

**WB (Western Blot)**

Western blot analysis of lysates from HepG2 cells, using Rev-Erbα/NR1D1 Rabbit pAb at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

ADDITIONAL INFORMATION

Related Product Information for anti-NR1D1 antibody	Background: This gene encodes a transcription factor that is a member of the nuclear receptor subfamily 1. The encoded protein is a ligand-sensitive transcription factor that negatively regulates the expression of core clock proteins. In particular this protein represses the circadian clock transcription factor aryl hydrocarbon receptor nuclear translocator-like protein 1 (ARNTL). This protein may also be involved in regulating genes that function in metabolic, inflammatory and cardiovascular processes.
Product Categories/Family for anti-NR1D1 antibody	Epigenetics & Nuclear Signaling ; Transcription Factors

NCBI AND UNIPROT INFORMATION

NCBI GI #	119100
NCBI GeneID	9572
UniProt Accession #	P20393 ; Q0P5Z4 ; Q15304
Molecular Weight	Observed MW: 78kDa calculated MW: 67kDa
NCBI Official Full Name	Nuclear receptor subfamily 1 group D member 1
NCBI Official Synonym Full Names	nuclear receptor subfamily 1, group D, member 1
NCBI Official Symbol	NR1D1
NCBI Official Synonym Symbols	EAR1; hRev; THRA1; THRAL; ear-1
NCBI Protein Information	nuclear receptor subfamily 1 group D member 1; Rev-ErbAalpha; rev-erbA-alpha; V-erbA-related protein 1; nuclear receptor Rev-ErbA-alpha
UniProt Protein Name	Nuclear receptor subfamily 1 group D member 1
UniProt Gene Name	NR1D1
UniProt Synonym Gene Names	EAR1; HREV; THRAL; EAR-1
UniProt Entry Name	NR1D1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rev-erbalpha-nr1d1-antibody-western-blot-immunohistochemistry-immunofluorescence-immunocytochemistry-elisa/28372>