

Rabbit anti-Human CYP2C19 Polyclonal Antibody | anti-CYP2C19 antibody

CYP2C19 Antibody (Center)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA284130	Rabbit	Human	Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

CYP2C19; Polyclonal Antibody; CYP2C19 Antibody (Center); Cytochrome P450 2C19; 11413-; (R)-limonene 6-monooxygenase; (S)-limonene 6-monooxygenase; (S)-limonene 7-monooxygenase; CYP11C17; CYP11C19; Cytochrome P450-11A; Cytochrome P450-254C; Mephenytoin 4-hydroxylase; anti-CYP2C19 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This CYP2C19 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 257-285 amino acids from the Central region of human CYP2C19.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	257-285
Sequence Length	490
Applicable Applications for anti-CYP2C19 antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
CYP2C19 Antibody (Center) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of CYP2C19 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



WB (Western Blot)
Western blot analysis of CYP2C19 Antibody (Center) in Jurkat cell line lysates (35ug/lane). CYP2C19 (arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-CYP2C19 antibody	CYP2C19 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, omeprazole, diazepam and some barbiturates. Polymorphism within its gene is associated with variable ability to metabolize mephenytoin, known as the poor metabolizer and extensive metabolizer phenotypes.
Product Categories/Family for anti-CYP2C19 antibody	Cardiovascular; Signal Transduction
References	Shuldiner,A.R., JAMA 302 (8), 849-857 (2009) Nelson,D.R., Pharmacogenetics 14 (1), 1-18 (2004)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503219
NCBI GeneID	1557
NCBI Accession #	NP_000760.1
NCBI GenBank Nucleotide #	NM_000769.2
UniProt Accession #	P33261; P33259; Q8WZB1; Q8WZB2; Q9UCD4
Molecular Weight	55931
NCBI Official Full Name	cytochrome P450 2C19
NCBI Official Synonym Full Names	cytochrome P450, family 2, subfamily C, polypeptide 19
NCBI Official Symbol	CYP2C19
NCBI Official Synonym Symbols	CPCJ; CYP2C; P450C2C; CYPIIC17; CYPIIC19; P450IIC19
NCBI Protein Information	cytochrome P450 2C19
UniProt Protein Name	Cytochrome P450 2C19
UniProt Gene Name	CYP2C19
UniProt Entry Name	CP2CJ_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyp2c19-antibody-immunohistochemistry-western-blot-elisa/284130>