

Rabbit PSME3 Antibody | anti-PSME3 antibody

PSME3 Rabbit pAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28429	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry


PRODUCT SYNONYMS

PSME3; PSME3 Rabbit pAb; HEL-S-283; Ki; PA28-gamma; PA28G; PA28gamma; REG-GAMMA; proteasome activator subunit 3; anti-PSME3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.09% Sodium azide, 50% glycerol, pH 7.3.
Sequence	RLSME
Applicable Applications for anti-PSME3 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry)
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200 IF/ICC: 1:50-1:200
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of human PSME3 (NP_005780.2).
Cellular Location	Cytoplasm, Nucleus
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of U-2 OS cells using PSME3 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of NIH-3T3 cells using PSME3 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of C6 cells using PSME3 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat kidney using [KO Validated] PSME3 Rabbit pAb at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse lung using [KO Validated] PSME3 Rabbit pAb at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human esophageal using [KO Validated] PSME3 Rabbit pAb at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

ADDITIONAL INFORMATION

Related Product Information for anti-PSME3 antibody	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. The immunoproteasome contains an alternate regulator, referred to as the 11S regulator or PA28, that replaces the 19S regulator. Three subunits (alpha, beta and gamma) of the 11S regulator have been identified. This gene encodes the gamma subunit of the 11S regulator. Six gamma subunits combine to form a homohexameric ring. Alternate splicing results in multiple transcript variants.
Product Categories/Family for anti-PSME3 antibody	Cancer; Tumor suppressors; p53 pathway; Cell Biology & Developmental Biology; Cell Cycle; Cell cycle inhibitors; Ubiquitin; Immunology & Inflammation; Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557795
NCBI GeneID	4771
NCBI Accession #	NP_000259.1
NCBI GenBank Nucleotide #	NM_000268.3

UniProt Accession #	P35240; O95683; Q8WUJ2; Q969N0; Q969Q3; Q96T30; Q96T31; Q96T32; Q96T33; Q9BTW3; Q9UNG9; Q9UNH3
Molecular Weight	69,690 Da
NCBI Official Full Name	merlin isoform 1
NCBI Official Synonym Full Names	neurofibromin 2 (merlin)
NCBI Official Symbol	NF2
NCBI Official Synonym Symbols	ACN; SCH; BANF
NCBI Protein Information	merlin; schwannomin; schwannomerlin; neurofibromin-2; moesin-ezrin-radixin like; moesin-ezrin-radixin-like protein; moesin-ezrin-radixin-like protein; neurofibromin 2 (bilateral acoustic neuroma)
UniProt Protein Name	Merlin
UniProt Gene Name	NF2
UniProt Synonym Gene Names	SCH
UniProt Entry Name	MERL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-human-mouse-rat-antibody/psme3/28429>