

Rabbit EphA2 Polyclonal Antibody | anti-EPHA2 antibody

EphA2 Antibody (C-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA284622	Rabbit	Human, hamster, mouse (Predicted Reactivity: Monkey)	ELISA, Western Blot


PRODUCT SYNONYMS

EphA2; Polyclonal Antibody; EphA2 Antibody (C-term); Ephrin type-A receptor 2; Epithelial cell kinase; Tyrosine-protein kinase receptor ECK; EPHA2; ECK; anti-EPHA2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, hamster, mouse (Predicted Reactivity: Monkey)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This EphA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 919-950 amino acids from the C-terminal region of human EphA2.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 1.17 (varies by lot)
Sequence Positions	919-950
Sequence Length	976
Applicable Applications for anti-EPHA2 antibody	ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)
Western blot analysis of EphA2 (arrow) using rabbit polyclonal EphA2 Antibody (C-term). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the EphA2 gene (Lane 2) (Origene Technologies).

ADDITIONAL INFORMATION

Related Product Information for anti-EPHA2 antibody	Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine kinase (TK) group is mainly involved in the regulation of cell-cell interactions such as differentiation, adhesion, motility and death. There are currently about 90 TK genes sequenced, 58 are of receptor protein TK (e.g. EGFR, EPH, FGFR, PDGFR, TRK, and VEGFR families), and 32 of cytosolic TK (e.g. ABL, FAK, JAK, and SRC families).
Product Categories/Family for anti-EPHA2 antibody	Cardiovascular; Neuroscience; Signal Transduction
References	Lindberg, R.A., et al., Mol. Cell. Biol. 10(12):6316-6324 (1990).

NCBI AND UNIPROT INFORMATION

NCBI GI #	32967311
NCBI GeneID	1969
NCBI Accession #	NP_004422.2
NCBI GenBank Nucleotide #	NM_004431.3
UniProt Accession #	P29317; Q8N3Z2; B5A968
Molecular Weight	108266
NCBI Official Full Name	ephrin type-A receptor 2
NCBI Official Synonym Full Names	EPH receptor A2
NCBI Official Symbol	EPHA2
NCBI Official Synonym Symbols	ECK; CTPA; ARCC2; CTPP1; CTRCT6
NCBI Protein Information	ephrin type-A receptor 2
UniProt Protein Name	Ephrin type-A receptor 2
UniProt Gene Name	EPHA2

UniProt Synonym Gene Names	ECK
UniProt Entry Name	EPHA2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/epha2-antibody-western-blot-elisa/284622>