

Rabbit anti-Human IRF8 Polyclonal Antibody | anti-IRF8 antibody

IRF8 Antibody (C-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA284934	Rabbit	Human	Western Blot, ELISA, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

IRF8; Polyclonal Antibody; IRF8 Antibody (C-term); Interferon regulatory factor 8; IRF-8; Interferon consensus sequence-binding protein; H-ICSBP; ICSBP; ICSBP1; anti-IRF8 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This IRF8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 384-409 amino acids from the C-terminal region of human IRF8.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	384-409
Sequence Length	426
Applicable Applications for anti-IRF8 antibody	WB (Western Blot), ELISA, FCM/FACS (Flow Cytometry)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



FCM/FACS (Flow Cytometry)
IRF8 Antibody (C-term) flow cytometric analysis of HL-60 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



WB (Western Blot)
Western blot analysis of anti-IRF8 Antibody (C-term) in HL60 cell line lysates (35ug/lane). IRF8 (arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-IRF8 antibody	Interferon consensus sequence-binding protein (ICSBP) is a transcription factor of the interferon (IFN) regulatory factor (IRF) family. Proteins of this family are composed of a conserved DNA-binding domain in the N-terminal region and a divergent C-terminal region that serves as the regulatory domain. The IRF family proteins bind to the IFN-stimulated response element (ISRE) and regulate expression of genes stimulated by type I IFNs, namely IFN- α and IFN- β . IRF family proteins also control expression of IFN- α and IFN- β -regulated genes that are induced by viral infection.
Product Categories/Family for anti-IRF8 antibody	Immunology
References	McGough, J.M., Mol. Cancer Res. 6 (12), 1841-1851 (2008) Tshuikina, M., Exp. Hematol. 36 (12), 1673-1681 (2008) Martinez, A., Am. J. Surg. Pathol. 32 (8), 1190-1200 (2008)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504567
NCBI GeneID	3394
NCBI Accession #	NP_002154.1
NCBI GenBank Nucleotide #	NM_002163.2
UniProt Accession #	Q02556; A0AV82
Molecular Weight	48356
NCBI Official Full Name	interferon regulatory factor 8
NCBI Official Synonym Full Names	interferon regulatory factor 8
NCBI Official Symbol	IRF8
NCBI Official Synonym Symbols	ICSBP; IRF-8; ICSBP1; IMD32A; IMD32B; H-ICSBP
NCBI Protein Information	interferon regulatory factor 8
UniProt Protein Name	Interferon regulatory factor 8
UniProt Gene Name	IRF8
UniProt Synonym Gene Names	ICSBP1; IRF-8; H-ICSBP; ICSBP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/irf8-antibody-western-blot-flow-cytometry-elisa/284934>