

Rabbit anti-Mouse Trem2 Polyclonal Antibody | anti-Trem2 antibody

Mouse Trem2 Antibody (N-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA285225	Rabbit	Mouse	Immunohistochemistry, ELISA, Western Blot

PRODUCT SYNONYMS


Trem2; Polyclonal Antibody; Mouse Trem2 Antibody (N-term); Triggering receptor expressed on myeloid cells 2; TREM-2; Triggering receptor expressed on monocytes 2; Trem2a; Trem2b; Trem2c; anti-Trem2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	This Mouse Trem2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 22-48 amino acids from the N-terminal region of mouse Trem2.
Purity/Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Sequence Positions	22-48
Sequence Length	249
Applicable Applications for anti-Trem2 antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Function	Forms a receptor signaling complex with TYROBP which mediates signaling and cell activation following ligand binding (PubMed:11241283). Acts as a receptor for amyloid-beta protein 42, a cleavage product of the amyloid-beta precursor protein APP, and mediates its uptake and degradation by microglia (PubMed:27477018, PubMed:29518356). Binding to amyloid-beta 42 mediates microglial activation, proliferation, migration, apoptosis and expression of proinflammatory cytokines, such as IL6R and CCL3, and the anti-inflammatory cytokine ARG1 (PubMed:27477018, PubMed:29518356). Acts as a receptor for lipoprotein particles such as LDL, VLDL, and HDL and for apolipoproteins such as APOA1, APOA2, APOB, APOE, APOE2, APOE3, APOE4, and CLU and enhances their uptake in microglia (PubMed:27477018). Binds phospholipids (preferably anionic lipids) such as phosphatidylserine, phosphatidylethanolamine, phosphatidylglycerol and sphingomyelin (By similarity). Regulates microglial proliferation by acting as an upstream regulator of the Wnt/beta-catenin signaling cascade (PubMed:28077724). Required for microglial phagocytosis of apoptotic neurons (PubMed:24990881). Also required for microglial activation and phagocytosis of myelin debris after neuronal injury and of neuronal synapses during synapse elimination in the developing brain (PubMed:15728241, PubMed:25631124, PubMed:28592261, PubMed:29752066). Regulates microglial chemotaxis and process outgrowth, and also the microglial response to oxidative stress and lipopolysaccharide (PubMed:28483841, PubMed:29663649, PubMed:29859094, PubMed:30232263). It suppresses PI3K and NF-kappa-B signaling in response to lipopolysaccharide; thus promoting phagocytosis, suppressing proinflammatory cytokine and nitric oxide production, inhibiting apoptosis and increasing expression of IL10 and TGFB (PubMed:29663649). During oxidative stress, it promotes anti-apoptotic NF-kappa-B signaling and ERK signaling (PubMed:28592261). Plays a role in microglial MTOR activation and metabolism (PubMed:28802038). Regulates age-related changes in microglial numbers (PubMed:25631124, PubMed:29752066, PubMed:30548312). Triggers activation of the immune responses in macrophages and dendritic cells. Mediates cytokine-induced formation of multinucleated giant cells which are formed by the fusion of macrophages (PubMed:18957693). In dendritic cells, receptor of SEMA6D with PLEXNA1 as coreceptor and mediates up-regulation of chemokine receptor CCR7 and dendritic cell maturation and survival (PubMed:16715077). Involved in the positive regulation of osteoclast differentiation (PubMed:16418779).

Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Expressed in the brain, specifically in microglia (at protein level) (PubMed:15728241, PubMed:27477018, PubMed:28077724, PubMed:28559417, PubMed:28592261, PubMed:28802038, PubMed:28855301, PubMed:29752066, PubMed:29794134). Expressed in macrophages (at protein level) (PubMed:11241283, PubMed:28559417, PubMed:28802038). Expressed at higher levels in the CNS, heart and lung than in lymph nodes or in other non-lymphoid tissues such as kidney, liver and testis (PubMed:12472885). In the CNS not all microglia express TREM2 (PubMed:12472885). Brain regions with an incomplete blood-brain barrier had the lowest percentages of TREM2 expressing microglia, whereas the lateral entorhinal and cingulate cortex had the highest percentages (PubMed:12472885).
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES




IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded mouse lung tissue using AAA285225 performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



IHC (Immunohistochemisry)
Immunohistochemical analysis of paraffin-embedded mouse brain tissue using AAA285225 performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



WB (Western Blot)
All lanes: Anti-Mouse Trem2 Antibody (N-term) at 1:500 dilutionLane 1: Mouse lung tissue lysateLane 2: Mouse liver tissue lysate Lysates/proteins at 20 ug per lane.Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.Predicted band size : 25 kDaBlocking/Dilution buffer: 5% NFDm/TBST.



WB (Western Blot)
All lanes: Anti-Mouse Trem2 Antibody (N-term) at 1:500-1000 dilutionLane 1: Mouse kidney tissue lysateLane 2: Mouse testis tissue lysate Lysates/proteins at 20 ug per lane.Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.Predicted band size: 25 kDaBlocking/Dilution buffer: 5% NFDm/TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-Trem2 antibody	Trem2 may have a role in chronic inflammations and may stimulate production of constitutive rather than inflammatory chemokines and cytokines. Forms a receptor signaling complex with TYROBP and triggers activation of the immune responses in macrophages and dendritic cells.
---	---

Product Categories/Family for anti-Trem2 antibody	Immunology
References	Koth, L.L., et al. J. Immunol. 184(11):6522-6528(2010) Whittaker, G.C., et al. J. Biol. Chem. 285(5):2976-2985(2010) Peng, Q., et al. Sci Signal 3 (122), RA38 (2010) : Chang, J.H., et al. Biochem. Biophys. Res. Commun. 389(1):28-33(2009) Hsieh, C.L., et al. J. Neurochem. 109(4):1144-1156(2009)

NCBI AND UNIPROT INFORMATION

NCBI GI #	440918700
NCBI GeneID	83433
NCBI Accession #	NP_001259007.1 ; NP_112544.1
NCBI GenBank Nucleotide #	NM_001272078.1
UniProt Accession #	Q99NH8 ; Q8CGK4 ; Q8CIA6 ; Q99NH9 ; Q9JL34
NCBI Official Full Name	triggering receptor expressed on myeloid cells 2 isoform 2
NCBI Official Synonym Full Names	triggering receptor expressed on myeloid cells 2
NCBI Official Symbol	Trem2
NCBI Official Synonym Symbols	TREM-2; Trem2a; Trem2b; Trem2c
NCBI Protein Information	triggering receptor expressed on myeloid cells 2
UniProt Protein Name	Triggering receptor expressed on myeloid cells 2
UniProt Gene Name	Trem2
UniProt Synonym Gene Names	Trem2a; Trem2b; Trem2c; TREM-2
UniProt Entry Name	TREM2_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/trem2-antibody-immunohistochemistry-western-blot-elisa/285225>