

Rabbit ATG7 Polyclonal Antibody | anti-ATG7 antibody

ATG7 Antibody (Center)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA285513	Rabbit	Human, mouse (Predicted Reactivity: Chicken, Rat)	Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

ATG7; Polyclonal Antibody; ATG7 Antibody (Center); Ubiquitin-like modifier-activating enzyme ATG7; ATG12-activating enzyme E1 ATG7; Autophagy-related protein 7; APG7-like; hAGP7; Ubiquitin-activating enzyme E1-like protein; APG7L; anti-ATG7 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, mouse (Predicted Reactivity: Chicken, Rat)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This ATG7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 284-313 amino acids from the Central region of human ATG7.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	284-313
Sequence Length	623
Applicable Applications for anti-ATG7 antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES



IHC (Immunohistochemistry)
Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



WB (Western Blot)
The ATG7 Antibody (Center) Pab is used in Western blot to detect APG7L in mouse liver tissue lysate. APG7L (arrow) was detected using the purified Pab.



WB (Western Blot)
Western blot analysis of anti-ATG7 Antibody (Center) Pab in 293 cell line lysates transiently transfected with the ATG7 gene (2ug/lane). hAPG7L-P299(arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-ATG7 antibody	Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). APG7 functions as an E1 enzyme essential for multisubstrates such as GABARAPL1 and ATG12. APG3L is an E2-like conjugating enzyme facilitating covalent binding of APG8 (MAP1LC3) to phosphatidylethanolamine (PE). APG7 (an E1-like enzyme) facilitates this reaction by forming an E1-E2 complex with APG3. Formation of the PE conjugate is essential for autophagy.
Product Categories/Family for anti-ATG7 antibody	Cancer; Cardiovascular; Cell Biology; Metabolism; Signal Transduction; Autophagy
References	Baehrecke EH. Nat Rev Mol Cell Biol. 6(6):505-10. (2005) Lum JJ, et al. Nat Rev Mol Cell Biol. 6(6):439-48. (2005) Greenberg JT. Dev Cell. 8(6):799-801. (2005) Levine B. Cell. 120(2):159-62. (2005) Shintani T and Klionsky DJ. Science. 306(5698):990-5. (2004) Tanida I., et al. Biochem. Biophys. Res. Commun. 292:256-262(2002) Tanida I., et al. J. Biol. Chem. 277:13739-13744(2002)

NCBI AND UNIPROT INFORMATION

NCBI GI #	222144227
-----------	-----------

NCBI GeneID	10533
NCBI Accession #	NP_001129503.2; NP_001138384.1; NP_006386.1
NCBI GenBank Nucleotide #	NM_001136031.2
UniProt Accession #	O95352; Q7L8L0; Q9BWP2; Q9UFH4; B4E170; E9PB95
Molecular Weight	77960
NCBI Official Full Name	ubiquitin-like modifier-activating enzyme ATG7 isoform b
NCBI Official Synonym Full Names	autophagy related 7
NCBI Official Symbol	ATG7
NCBI Official Synonym Symbols	GSA7; APG7L; APG7-LIKE
NCBI Protein Information	ubiquitin-like modifier-activating enzyme ATG7
UniProt Protein Name	Ubiquitin-like modifier-activating enzyme ATG7
UniProt Gene Name	ATG7
UniProt Synonym Gene Names	APG7L; APG7-like; hAGP7
UniProt Entry Name	ATG7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/atg7-antibody-immunohistochemistry-western-blot-elisa/285513>