

Rabbit ILK Polyclonal Antibody | anti-ILK antibody

ILK Antibody (T173)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA285816	Rabbit	Human (Predicted Reactivity: Bovine, Mouse, Rat)	Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

ILK; Polyclonal Antibody; ILK Antibody (T173); Integrin-linked protein kinase; 59 kDa serine/threonine-protein kinase; ILK-1; ILK-2; p59ILK; ILK1; ILK2; anti-ILK antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human (Predicted Reactivity: Bovine, Mouse, Rat)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This ILK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 151-180 amino acids from human ILK.
Purity/Purification	Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Concentration	Vial Concentration: 0.5 (varies by lot)
Sequence Positions	151-180
Sequence Length	452
Applicable Applications for anti-ILK antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
Formalin-fixed and paraffin-embedded human skeletal muscle tissue reacted with ILK Antibody (T173) antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)
Western blot analysis of anti-ILK Antibody (T173) (RB13894) in NCI-H460 cell line lysates (35ug/lane). EN1(arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-ILK antibody	Transduction of extracellular matrix signals through integrins influences intracellular and extracellular functions, and appears to require interaction of integrin cytoplasmic domains with cellular proteins. Integrin-linked kinase (ILK), interacts with the cytoplasmic domain of beta-1 integrin. ILK is a serine/threonine protein kinase with 4 ankyrin-like repeats, which associates with the cytoplasmic domain of beta integrins and acts as a proximal receptor kinase regulating integrin-mediated signal transduction.
Product Categories/Family for anti-ILK antibody	Signal Transduction
References	Li, Y., et al., J. Clin. Invest. 112(4):503-516 (2003). Troussard, A.A., et al., J. Biol. Chem. 278(25):22374-22378 (2003). Marotta, A., et al., Br. J. Cancer 88(11):1755-1762 (2003). Cordes, N., et al., Br. J. Cancer 88(9):1470-1479 (2003). Fukuda, T., et al., J. Cell Biol. 160(7):1001-1008 (2003).

NCBI AND UNIPROT INFORMATION

NCBI GI #	62420873
NCBI GeneID	3611
NCBI Accession #	NP_001014794.1; NP_001014795.1; NP_001265370.1; NP_004508.1
NCBI GenBank Nucleotide #	NM_001014794.2
UniProt Accession #	Q13418; P57043; Q68DZ3; B7Z1I0; B7Z418; D3DQU0
Molecular Weight	51419
NCBI Official Full Name	integrin-linked protein kinase isoform 1
NCBI Official Synonym Full Names	integrin-linked kinase
NCBI Official Symbol	ILK
NCBI Official Synonym Symbols	P59; ILK-1; ILK-2; p59ILK; HEL-S-28
NCBI Protein Information	integrin-linked protein kinase
UniProt Protein Name	Integrin-linked protein kinase
UniProt Gene Name	ILK

UniProt Synonym Gene Names	ILK1; ILK2
UniProt Entry Name	ILK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ilk-antibody-immunohistochemistry-western-blot-elisa/285816>