

Rabbit BPTF Polyclonal Antibody | anti-BPTF antibody

BPTF Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28813	Rabbit	Rat, Dog, Cow	Immunohistochemistry, Immunohistochemistry, Immunofluorescence, Immunocytochemistry


PRODUCT SYNONYMS

BPTF; Polyclonal Antibody; BPTF Polyclonal Antibody; Nucleosome-remodeling factor subunit BPTF; Bromodomain and PHD finger-containing transcription factor; Fetal Alz-50 clone 1 protein; Fetal Alzheimer antigen; FAC1; FALZ; anti-BPTF antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-BPTF antibody	IHC (Immunohistochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry-Paraffin)

Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (BPTF) Polyclonal Antibody, Unconjugated (bs-11641R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




IHC (Immunohistochemistry-Paraffin)

Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (BPTF) Polyclonal Antibody, Unconjugated (bs-11641R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (BPTF) Polyclonal Antibody, Unconjugated (bs-11641R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry-Paraffin)

Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min; Incubation: Anti-BPTF Polyclonal Antibody, Unconjugated(bs-11641R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

**IHC (Immunohistochemistry-Paraffin)**

Dilution: Tissue/cell: rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min; Incubation: Anti-BPTF Polyclonal Antibody, Unconjugated(bs-11641R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

**IHC (Immunohistochemistry-Paraffin)**

Tissue/cell: rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min; Incubation: Anti-BPTF Polyclonal Antibody, Unconjugated(bs-11641R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	38788260
NCBI GeneID	2186
NCBI Accession #	NP_004450.3
NCBI GenBank Nucleotide #	NM_004459.6
UniProt Accession #	Q12830; Q6NX67; Q7Z7D6; Q9UIG2
Molecular Weight	322,217 Da
NCBI Official Full Name	nucleosome-remodeling factor subunit BPTF isoform 2
NCBI Official Synonym Full Names	bromodomain PHD finger transcription factor
NCBI Official Symbol	BPTF
NCBI Official Synonym Symbols	FAC1; FALZ; NURF301
NCBI Protein Information	nucleosome-remodeling factor subunit BPTF; fetal Alzheimer antigen; fetal Alz-50 clone 1 protein; fetal Alz-50 reactive clone 1; nucleosome remodeling factor, large subunit; bromodomain and PHD domain transcription factor; bromodomain and PHD finger-conta
UniProt Protein Name	Nucleosome-remodeling factor subunit BPTF
UniProt Gene Name	BPTF
UniProt Synonym Gene Names	FAC1; FALZ
UniProt Entry Name	BPTF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/bptf-antibody-immunohistochemistry-immunohistochemistry-immunofluorescence-immunocytochemistry-elisa/28813>