

Rabbit CDC23 Polyclonal Antibody | anti-CDC23 antibody

CDC23 Antibody (N-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA288643	Rabbit	Human (Predicted Reactivity: Bovine)	Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

CDC23; Polyclonal Antibody; CDC23 Antibody (N-term); Cell division cycle protein 23 homolog; Anaphase-promoting complex subunit 8; APC8; Cyclosome subunit 8; ANAPC8; anti-CDC23 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human (Predicted Reactivity: Bovine)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This CDC23 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 140-168 amino acids from the N-terminal region of human CDC23.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	140-168
Sequence Length	597
Applicable Applications for anti-CDC23 antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed and paraffin-embedded human brain tissue reacted with CDC23 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)
Western blot analysis of CDC23 antibody (N-term) in HL60 cell line lysates(35ug/lane). CDC23 (arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-CDC23 antibody	CDC23 shares strong similarity with Saccharomyces cerevisiae Cdc23, a protein essential for cell cycle progression through the G2/M transition. This protein is a component of anaphase-promoting complex (APC), which is composed of eight protein subunits and highly conserved in eukaryotic cells. APC catalyzes the formation of cyclin B-ubiquitin conjugate that is responsible for the ubiquitin-mediated proteolysis of B-type cyclins. This protein and 3 other members of the APC complex contain the TPR (tetratricopeptide repeat), a protein domain important for protein-protein interaction.
Product Categories/Family for anti-CDC23 antibody	Cancer; Cell biology
References	Zhao,N., Genomics 53 (2), 184-190 (1998)

NCBI AND UNIPROT INFORMATION

NCBI GI #	118402596
NCBI GeneID	8697
NCBI Accession #	NP_004652.2
NCBI GenBank Nucleotide #	NM_004661.3
UniProt Accession #	Q9UJX2; O75433; Q53FN2; Q9BS73; A8K6E5; B4E3A2; B7WP05; D3DQB7
Molecular Weight	68834
NCBI Official Full Name	cell division cycle protein 23 homolog
NCBI Official Synonym Full Names	cell division cycle 23
NCBI Official Symbol	CDC23
NCBI Official Synonym Symbols	APC8; CUT23; ANAPC8
NCBI Protein Information	cell division cycle protein 23 homolog
UniProt Protein Name	Cell division cycle protein 23 homolog
UniProt Gene Name	CDC23
UniProt Synonym Gene Names	ANAPC8; APC8
UniProt Entry Name	CDC23_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cdc23-antibody-immunohistochemistry-western-blot-elisa/288643>