

Rabbit BAI1 Polyclonal Antibody | anti-BAI1 antibody

BAI1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28895	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

BAI1; Polyclonal Antibody; BAI1 Polyclonal Antibody; Brain-specific angiogenesis inhibitor 1; anti-BAI1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-BAI1 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



Application Data

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.




Application Data

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



Application Data

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-BAI1 antibody

Phosphatidylserine receptor which enhances the engulfment of apoptotic cells (PubMed:24509909). Also mediates the binding and engulfment of Gram-negative bacteria (PubMed:26838550). Stimulates production of reactive oxygen species by macrophages in response to Gram-negative bacteria, resulting in enhanced microbicidal macrophage activity (PubMed:26838550). In the gastric mucosa, required for recognition and engulfment of apoptotic gastric epithelial cells (PubMed:24509909). Promotes myoblast fusion (By similarity). Activates the Rho pathway in a G-protein-dependent manner (PubMed:23782696). Inhibits MDM2-mediated ubiquitination and degradation of DLG4/PSD95, promoting DLG4 stability and regulating synaptic plasticity (By similarity). Required for the formation of dendritic spines by ensuring the correct localization of PARD3 and TIAM1 (By similarity). Potent inhibitor of angiogenesis in brain and may play a significant role as a mediator of the p53/TP53 signal in suppression of glioblastoma (PubMed:11875720).

NCBI AND UNIPROT INFORMATION

NCBI GI #	111118994
NCBI GeneID	575
NCBI Accession #	NP_001693.2
NCBI GenBank Nucleotide #	NM_001702.2
UniProt Accession #	O14514
Molecular Weight	173,501 Da
NCBI Official Full Name	brain-specific angiogenesis inhibitor 1
NCBI Official Synonym Full Names	brain-specific angiogenesis inhibitor 1

NCBI Official Symbol	BAI1
NCBI Official Synonym Symbols	GDAIF
NCBI Protein Information	brain-specific angiogenesis inhibitor 1
UniProt Protein Name	Brain-specific angiogenesis inhibitor 1
UniProt Gene Name	BAI1
UniProt Entry Name	BAI1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/bai1-antibody-western-blot-immunohistochemistry-elisa/28895>