

Rabbit ENOA Polyclonal Antibody | anti-ENO1 antibody

ENOA Antibody (C-term)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA288994	Rabbit	Human, mouse (Predicted Reactivity: Monkey)	Western Blot, Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS


ENOA; Polyclonal Antibody; ENOA Antibody (C-term); Alpha-enolase; 2-phospho-D-glycerate hydro-lyase; C-myc promoter-binding protein; Enolase 1; MBP-1; MPB-1; Non-neural enolase; NNE; Phosphopyruvate hydratase; Plasminogen-binding protein; ENO1; ENO1L1; MBPB1; MPB1; anti-ENO1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, mouse (Predicted Reactivity: Monkey)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This ENOA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 405-434 amino acids from the C-terminal region of human ENOA.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	405-434
Sequence Length	341
Applicable Applications for anti-ENO1 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Crown Antibody	Yes
Function	Multifunctional enzyme that, as well as its role in glycolysis, plays a part in various processes such as growth control, hypoxia tolerance and allergic responses. May also function in the intravascular and pericellular fibrinolytic system due to its ability to serve as a receptor and activator of plasminogen on the cell surface of several cell-types such as leukocytes and neurons. Stimulates immunoglobulin production.
Tissue Location	The alpha/alpha homodimer is expressed in embryo and in most adult tissues. The alpha/beta heterodimer and the beta/beta homodimer are found in striated muscle, and the alpha/gamma heterodimer and the gamma/gamma homodimer in neurons
Cellular Location	Cytoplasm. Cell membrane. Cytoplasm, myofibril, sarcomere, M line. Note=Can translocate to the plasma membrane in either the homodimeric (alpha/alpha) or heterodimeric (alpha/gamma) form. ENO1 is localized to the M line
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.




IHC (Immunohistochemistry)
ENOA Antibody (C-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ENOA Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



IF (Immunofluorescence)
Confocal immunofluorescent analysis of ENOA Antibody (C-term) with HeLa cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).



FCM/FACS (Flow Cytometry)
ENOA Antibody (C-term) flow cytometry analysis of Ramos cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



WB (Western Blot)
ENOA Antibody (C-term) western blot analysis in Jurkat, MCF-7, U251 cell line and mouse brain tissue lysates (35ug/lane). This demonstrates the ENOA antibody detected the ENOA protein (arrow).

ADDITIONAL INFORMATION

Related Product Information for anti-ENO1 antibody	ENO1 is one of three enolase isoenzymes found in mammals; the protein alpha-enolase, a homodimeric soluble enzyme, and is also a shorter monomeric structural lens protein, tau-crystallin. The two proteins are made from the same message. The full length protein, the isoenzyme, is found in the cytoplasm. The shorter protein is produced from an alternative translation start, is localized to the nucleus, and has been found to bind to an element in the c-myc promoter.
Product Categories/Family for anti-ENO1 antibody	Crown Antibodies; Cancer; Metabolism; Signal Transduction
References	Cappello,P., Int. J. Cancer 125 (3), 639-648 (2009) Wygrecka,M., Blood 113 (22), 5588-5598 (2009)

NCBI AND UNIPROT INFORMATION

NCBI GI #	319996655
NCBI GeneID	2023
NCBI Accession #	Q4R5L2; NP_001419.1
NCBI GenBank Nucleotide #	NM_001201483.1

UniProt Accession #	P06733; P22712; Q16704; Q4TUS4; Q53FT9; Q53HR3; Q658M5; Q6GMP2; Q71V37; Q7Z3V6; Q8WU71; B2RD59
Molecular Weight	47169
NCBI Official Full Name	c-myc promoter-binding protein-1 isoform MBP-1
NCBI Official Synonym Full Names	enolase 1, (alpha)
NCBI Official Symbol	ENO1
NCBI Official Synonym Symbols	NNE; PPH; MPB1; ENO1L1
NCBI Protein Information	alpha-enolase; c-myc promoter-binding protein-1
UniProt Protein Name	Alpha-enolase
UniProt Gene Name	ENO1
UniProt Synonym Gene Names	ENO1L1; MBPB1; MPB1; NNE
UniProt Entry Name	ENOA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/enoa-antibody-western-blot-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/288994>