

Rabbit anti-Human PRAK (MAPKAPK5) Polyclonal Antibody | anti-MAPKAPK5 antibody

PRAK (MAPKAPK5) Antibody (T182)

CATALOG # AAA289248	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

PRAK (MAPKAPK5); Polyclonal Antibody; PRAK (MAPKAPK5) Antibody (T182); MAP kinase-activated protein kinase 5; MAPK-activated protein kinase 5; MAPKAP kinase 5; MAPKAP-K5; MAPKAPK-5; MK-5; MK5; p38-regulated/activated protein kinase; PRAK; MAPKAPK5; anti-MAPKAPK5 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This PRAK(MAPKAPK5) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 160-189 amino acids from human PRAK(MAPKAPK5).
Purity/Purification	Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Concentration	Vial Concentration: 0.25 (varies by lot)
Sequence Positions	160-189
Sequence Length	471
Applicable Applications for anti-MAPKAPK5 antibody	ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)
Western blot analysis of MAPKAPK5 (arrow) using rabbit polyclonal MAPKAPK5 Antibody (T182). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the MAPKAPK5 gene.



WB (Western Blot)
Western blot analysis of MAPKAPK5 Antibody (T182) in Hela cell line lysates (35ug/lane). MAPKAPK5 (arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPKAPK5 antibody	MAPKAPK5 is a member of the serine/threonine kinase family. In response to cellular stress and proinflammatory cytokines, this kinase is activated through its phosphorylation by MAP kinases including MAPK1/ERK, MAPK14/p38-alpha, and MAPK11/p38-beta. In vitro, this kinase phosphorylates heat shock protein HSP27 at its physiologically relevant sites.
Product Categories/Family for anti-MAPKAPK5 antibody	Cancer
References	New, L., et al., EMBO J. 17(12):3372-3384 (1998). Ni, H., et al., Biochem. Biophys. Res. Commun. 243(2):492-496 (1998). Sudo T.,Biochem. Biophys. Res. Commun. 269:521-525(2000).

NCBI AND UNIPROT INFORMATION

NCBI GI #	21237765
NCBI GeneID	8550
NCBI Accession #	NP_003659.2; NP_620777.1
NCBI GenBank Nucleotide #	NM_003668.3
UniProt Accession #	Q8IW41; O60491; Q86X46; Q9BVX9; Q9UG86; B3KVA5
Molecular Weight	54220
NCBI Official Full Name	MAP kinase-activated protein kinase 5 isoform 1
NCBI Official Synonym Full Names	mitogen-activated protein kinase-activated protein kinase 5
NCBI Official Symbol	MAPKAPK5
NCBI Official Synonym Symbols	MK5; MK-5; PRAK; MAPKAP-K5
NCBI Protein Information	MAP kinase-activated protein kinase 5
UniProt Protein Name	MAP kinase-activated protein kinase 5
UniProt Gene Name	MAPKAPK5
UniProt Synonym Gene Names	PRAK; MAPK-activated protein kinase 5; MAPKAP kinase 5; MAPKAP-K5; MAPKAPK-5; MK-5; MK5; PRAK
UniProt Entry Name	MAPK5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/prak-mapkapk5-antibody-western-blot-elisa/289248>