

Rabbit anti-Human TrkC Polyclonal Antibody | anti-NTRK3 antibody

TrkC Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA289528	Rabbit	Human	Western Blot, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

TrkC; Polyclonal Antibody; TrkC Antibody; NT-3 growth factor receptor; GP145-TrkC; Trk-C; Neurotrophic tyrosine kinase receptor type 3; TrkC tyrosine kinase; NTRK3; TRKC; anti-NTRK3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This TrkC antibody is generated from rabbits immunized with a his tag recombinant protein of human TrkC.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Length	817
Applicable Applications for anti-NTRK3 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Antigen Type	Recombinant Protein
Crown Antibody	Yes
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)
TrkC Antibody western blot analysis in U87-MG cell line lysates (35ug/lane).This demonstrates the TrkC antibody detected the TrkC protein (arrow).



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded H. brain section using TrkC Antibody. AAA289528 was diluted at 1:100 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

ADDITIONAL INFORMATION

Related Product Information for anti-NTRK3 antibody	TRKC, a member of the insuline receptor subfamily of Tyr protein kinases, is a receptor for neurotrophin-3 (NT-3). Known substrates for the TRK receptors are SHC, PI-3 kinase, and PLCG1. The different isoforms do not have identical signaling properties. The protein is widely expressed, mainly in the nervous tissue. The isoform B is expressed in a relatively large amount in the adult brain comparatively to fetal brain. TRKC is subject to ligand-mediated auto-phosphorylation. The protein structure contains 2 immunoglobulin-like C2-type domains and 2 leucine-rich (LRR) repeats.
Product Categories/Family for anti-NTRK3 antibody	Crown Antibodies; Cancer; Neuroscience; Signal Transduction
References	McGregor, L.M., et al., Genomics 22(2):267-272 (1994). Shelton, D.L., et al., J. Neurosci. 15 (1 Pt 2), 477-491 (1995).

NCBI AND UNIPROT INFORMATION

NCBI GI #	59889562
NCBI GeneID	4916
NCBI Accession #	NP_001012338.1; NP_002521.2
NCBI GenBank Nucleotide #	NM_001012338.2
UniProt Accession #	Q16288; O75682; Q12827; Q16289; B7Z4C5; E9PG56; H0YND1
Molecular Weight	94428
NCBI Official Full Name	NT-3 growth factor receptor isoform a
NCBI Official Synonym Full Names	neurotrophic tyrosine kinase, receptor, type 3
NCBI Official Symbol	NTRK3
NCBI Official Synonym Symbols	TRKC; GP145-TrkC; gp145(trkC)
NCBI Protein Information	NT-3 growth factor receptor
UniProt Protein Name	NT-3 growth factor receptor
UniProt Gene Name	NTRK3
UniProt Synonym Gene Names	TRKC; Trk-C

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/trkc-antibody-western-blot-immunohistochemistry-elisa/289528>