

Rabbit PERK Polyclonal Antibody | anti-PERK antibody

PERK (phospho Thr981) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28957	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

PERK; Polyclonal Antibody; PERK (phospho Thr981) Polyclonal Antibody; EIF2AK3; PEK; Eukaryotic translation initiation factor 2-alpha kinase 3; PRKR-like endoplasmic reticulum kinase; Pancreatic eIF2-alpha kinase; HsPEK; anti-PERK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Applicable Applications for anti-PERK antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Application Notes	IF: 1:50-200 IHC: 1/100-1/300 ELISA: 1/40000
Cellular Location	Endoplasmic reticulum membrane; Single-pass type I membrane protein
Tissue Location	Ubiquitous. A high level expression is seen in secretory tissues
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: IF: 1:50-200 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: IF: 1:50-200 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: IF: 1:50-200 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-PERK antibody

Metabolic-stress sensing protein kinase that phosphorylates the alpha subunit of eukaryotic translation initiation factor 2 (eIF-2-alpha/EIF2S1) on 'Ser-52' during the unfolded protein response (UPR) and in response to low amino acid availability. Converts phosphorylated eIF-2-alpha/EIF2S1 either in a global protein synthesis inhibitor, leading to a reduced overall utilization of amino acids, or to a translation initiation activator of specific mRNAs, such as the transcriptional activator ATF4, and hence allowing ATF4-mediated reprogramming of amino acid biosynthetic gene expression to alleviate nutrient depletion. Serves as a critical effector of unfolded protein response (UPR)- induced G1 growth arrest due to the loss of cyclin-D1 (CCND1). Involved in control of mitochondrial morphology and function.

NCBI AND UNIPROT INFORMATION

NCBI GI #	134304838
NCBI GeneID	9451
NCBI Accession #	NP_004827.4
NCBI GenBank Nucleotide #	NM_004836.5
UniProt Accession #	Q9NZJ5 ; O95846 ; Q53QY0 ; Q53SB1 ; A0AVH1 ; A0AVH2 ; B2RCU9
Molecular Weight	125,216 Da
NCBI Official Full Name	eukaryotic translation initiation factor 2-alpha kinase 3
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 2-alpha kinase 3
NCBI Official Symbol	EIF2AK3
NCBI Official Synonym Symbols	PEK; WRS; PERK
NCBI Protein Information	eukaryotic translation initiation factor 2-alpha kinase 3; hsPEK; pancreatic EIF2-alpha kinase; PRKR-like endoplasmic reticulum kinase; eukaryotic translation initiation factor 2 alpha kinase 3
UniProt Protein Name	Eukaryotic translation initiation factor 2-alpha kinase 3
UniProt Gene Name	EIF2AK3

UniProt Synonym Gene Names	PEK; PERK; HsPEK
UniProt Entry Name	E2AK3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/perk-antibody-western-blot-immunohistochemistry-immunofluorescence-elisa/28957>