

Rabbit anti-Human CD79a Polyclonal Antibody | anti-CD79a antibody

CD79a Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28987	Rabbit	Human	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

CD79a; Polyclonal Antibody; CD79a Polyclonal Antibody; CD79A; IGA; MB1; B-cell antigen receptor complex-associated protein alpha chain; Ig-alpha; MB-1 membrane glycoprotein; Membrane-bound immunoglobulin-associated protein; Surface IgM-associated protein; CD antigen CD79a; anti-CD79a antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD79a antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: IHC-p: 100-300. Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: IHC-p: 100-300. Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: IHC-p: 100-300. Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: IHC-p: 100-300. Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: IHC-p: 100-300. Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: IHC-p: 100-300. Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CD79a antibody

Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src- family tyrosine kinases. Represses BCR signaling during development of immature B-cells.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502685
NCBI GeneID	973
NCBI Accession #	NP_001774.1
NCBI GenBank Nucleotide #	NM_001783.3
UniProt Accession #	P11912; Q53FB8; A0N775
Molecular Weight	25,038 Da
NCBI Official Full Name	B-cell antigen receptor complex-associated protein alpha chain isoform 1
NCBI Official Synonym Full Names	CD79a molecule, immunoglobulin-associated alpha
NCBI Official Symbol	CD79A
NCBI Official Synonym Symbols	IGA; MB-1
NCBI Protein Information	B-cell antigen receptor complex-associated protein alpha chain; ig-alpha; MB-1 membrane glycoprotein; surface IgM-associated protein; CD79a antigen (immunoglobulin-associated alpha); membrane-bound immunoglobulin-associated protein
UniProt Protein Name	B-cell antigen receptor complex-associated protein alpha chain
UniProt Gene Name	CD79A
UniProt Synonym Gene Names	IGA; MB1
UniProt Entry Name	CD79A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd79a-antibody-western-blot-immunohistochemistry-elisa/28987>