

Rabbit c-Myc Polyclonal Antibody | anti-c-Myc antibody

c-Myc Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28997	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

c-Myc; Polyclonal Antibody; c-Myc Polyclonal Antibody; MYC; BHLHE39; Myc proto-oncogene protein; Class E basic helix-loop-helix protein 39; bHLHe39; Proto-oncogene c-Myc; Transcription factor p64; anti-c-Myc antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-c-Myc antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.




IF (Immunofluorescence)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.





IF (Immunofluorescence)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.



IF (Immunofluorescence)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.



Application Data

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

Application Data

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-c-Myc antibody

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Activates the transcription of growth-related genes. Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed:24940000).

NCBI AND UNIPROT INFORMATION

NCBI GI #	71774083
NCBI GeneID	4609
NCBI Accession #	NP_002458.2
NCBI GenBank Nucleotide #	NM_002467.4
UniProt Accession #	P01106; P01107; Q14026; A8WFE7
Molecular Weight	48,804 Da
NCBI Official Full Name	myc proto-oncogene protein
NCBI Official Synonym Full Names	v-myc myelocytomatosis viral oncogene homolog (avian)
NCBI Official Symbol	MYC
NCBI Official Synonym Symbols	MRTL; c-Myc; bHLHe39
NCBI Protein Information	myc proto-oncogene protein; proto-oncogene c-Myc; transcription factor p64; class E basic helix-loop-helix protein 39; avian myelocytomatosis viral oncogene homolog; v-myc avian myelocytomatosis viral oncogene homolog; myc-related translation/localization
UniProt Protein Name	Myc proto-oncogene protein
UniProt Gene Name	MYC
UniProt Synonym Gene Names	BHLHE39; bHLHe39
UniProt Entry Name	MYC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/c-myc-antibody-western-blot-immunohistochemistry-immunofluorescence-elisa/28997>