

Rabbit anti-Human MUC20 Polyclonal Antibody | anti-MUC20 antibody

MUC20 Antibody (N-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA290025	Rabbit	Human	Immunofluorescence, ELISA, Western Blot


PRODUCT SYNONYMS

MUC20; Polyclonal Antibody; MUC20 Antibody (N-term); Mucin-20; MUC-20; KIAA1359; anti-MUC20 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This MUC20 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 61-90 amino acids from the N-terminal region of human MUC20.
Purity/Purification	Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	61-90
Sequence Length	674
Applicable Applications for anti-MUC20 antibody	IF (Immunofluorescence), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)
Confocal immunofluorescent analysis of MUC20 Antibody (N-term) with WiDr cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



WB (Western Blot)
Western blot analysis of MUC20 (arrow) using rabbit polyclonal MUC20 Antibody (N-term). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the MUC20 gene (Lane 2).

ADDITIONAL INFORMATION

Related Product Information for anti-MUC20 antibody	This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins secreted by many epithelial tissues to form an insoluble mucous barrier. The shorter isoform expressed by this gene is localized to the plasma membrane, whereas the longer isoform might be secreted. The C terminus of this protein associates with the multifunctional docking site of the met proto-oncogene and suppresses activation of some downstream met signaling cascades. The protein features a tandem repeat domain that varies between 2 and 6 copies in different individuals. Multiple transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-MUC20 antibody	Signal Transduction
References	Li,G., Am. J. Nephrol. 26 (1), 43-49 (2006) Suzuki,Y., Genome Res. 14 (9), 1711-1718 (2004) Higuchi,T., Mol. Cell. Biol. 24 (17), 7456-7468 (2004) Higuchi,T., J. Biol. Chem. 279 (3), 1968-1979 (2004)

NCBI AND UNIPROT INFORMATION

NCBI GI #	541444092
NCBI GeneID	200958
NCBI Accession #	NP_001269435.1
NCBI GenBank Nucleotide #	NM_001282506.1
UniProt Accession #	Q8N307; Q6UX97; Q76I83; Q76I85; Q86ST8; Q8NBY6; Q96KA1; Q9P2I8
Molecular Weight	71982
NCBI Official Full Name	mucin-20 isoform a
NCBI Official Synonym Full Names	mucin 20, cell surface associated
NCBI Official Symbol	MUC20
NCBI Official Synonym Symbols	MUC-20
NCBI Protein Information	mucin-20
UniProt Protein Name	Mucin-20
UniProt Gene Name	MUC20

UniProt Synonym Gene Names	KIAA1359; MUC-20
UniProt Entry Name	MUC20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/muc20-antibody-immunofluorescence-western-blot-elisa/290025>