

Rabbit anti-Human B2M Polyclonal Antibody | anti-B2M antibody

B2M Antibody (C-term)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA290201	Rabbit	Human	Western Blot, Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA

PRODUCT SYNONYMS


B2M; Polyclonal Antibody; B2M Antibody (C-term); Beta-2-microglobulin; Beta-2-microglobulin form pl 53; anti-B2M antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This B2M antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 86-118 amino acids from the C-terminal region of human B2M.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	86-118
Sequence Length	119
Applicable Applications for anti-B2M antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA
Antigen Type	Synthetic Peptide
Crown Antibody	Yes
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.



FCM/FACS (Flow Cytometry)
B2M Antibody (C-term) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



IHC (Immunohiostchemistry)
Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with B2M antibody (C-term) , which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)
B2M Antibody (C-term) western blot analysis in Jurkat cell line lysates (35ug/lane).This demonstrates the B2M antibody detected the B2M protein (arrow).

ADDITIONAL INFORMATION

Related Product Information for anti-B2M antibody	Beta-2-microglobulin is a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells.
Product Categories/Family for anti-B2M antibody	Crown Antibodies; Cancer; Cardiovascular; Metabolism
References	Sakata,M., J. Mol. Biol. 382 (5), 1242-1255 (2008) Huang,W.C., Clin. Cancer Res. 14 (17), 5341-5347 (2008) Gattoni-Celli,S., Cancer Res. 52 (5), 1201-1204 (1992)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757826
NCBI GeneID	567
NCBI Accession #	NP_004039.1
NCBI GenBank Nucleotide #	NM_004048.2
UniProt Accession #	P61769; P01884; Q540F8; Q6IAT8; Q9UCK0; Q9UD48; Q9UDF4
Molecular Weight	13715

NCBI Official Full Name	beta-2-microglobulin
NCBI Official Synonym Full Names	beta-2-microglobulin
NCBI Official Symbol	B2M
NCBI Protein Information	beta-2-microglobulin
UniProt Protein Name	Beta-2-microglobulin
UniProt Gene Name	B2M
UniProt Entry Name	B2MG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/b2m-antibody-western-blot-flow-cytometry-immunohistochemistry-elisa/290201>