

Rabbit anti-Human Ki-67 Polyclonal Antibody | anti-Ki-67 antibody

Ki-67 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29057	Rabbit	Human	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

Ki-67; Polyclonal Antibody; Ki-67 Polyclonal Antibody; MKI67; Antigen KI-67; anti-Ki-67 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-Ki-67 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.




IF (Immunofluorescence)

Dilution: Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.




Application Data

Dilution: Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.




IF (Immunofluorescence)

Dilution: Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.



IF (Immunofluorescence)

Dilution: Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.



Application Data

Dilution: Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-Ki-67 antibody	Required to maintain individual mitotic chromosomes dispersed in the cytoplasm following nuclear envelope disassembly (PubMed:27362226). Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the chromosome surface (PubMed:27362226). Prevents chromosomes from collapsing into a single chromatin mass by forming a steric and electrostatic charge barrier: the protein has a high net electrical charge and acts as a surfactant, dispersing chromosomes and enabling independent chromosome motility (PubMed:27362226). Binds DNA, with a preference for supercoiled DNA and AT-rich DNA (PubMed:10878551). Does not contribute to the internal structure of mitotic chromosomes (By similarity). May play a role in chromatin organization (PubMed:24867636). It is however unclear whether it plays a direct role in chromatin organization or whether it is an indirect consequence of its function in maintaining mitotic chromosomes dispersed (Probable).
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	225543215
NCBI GeneID	4288
NCBI Accession #	NP_001139438.1
NCBI GenBank Nucleotide #	NM_001145966.1
UniProt Accession #	P46013; Q5VWH2
Molecular Weight	358,694 Da
NCBI Official Full Name	antigen KI-67 isoform 2
NCBI Official Synonym Full Names	antigen identified by monoclonal antibody Ki-67
NCBI Official Symbol	MKI67
NCBI Official Synonym Symbols	KIA
NCBI Protein Information	antigen KI-67; proliferation-related Ki-67 antigen
UniProt Protein Name	Antigen KI-67
UniProt Gene Name	MKI67
UniProt Entry Name	KI67_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.