

Rabbit anti-Human MMP-13 Polyclonal Antibody | anti-MMP-13 antibody

MMP-13 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29070	Rabbit	Human	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

MMP-13; Polyclonal Antibody; MMP-13 Polyclonal Antibody; MMP13; Collagenase 3; Matrix metalloproteinase-13; anti-MMP-13 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-MMP-13 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)


Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MMP-13 antibody

Plays a role in the degradation of extracellular matrix proteins including fibrillar collagen, fibronectin, TNC and ACAN. Cleaves triple helical collagens, including type I, type II and type III collagen, but has the highest activity with soluble type II collagen. Can also degrade collagen type IV, type XIV and type X. May also function by activating or degrading key regulatory proteins, such as TGFB1 and CTGF. Plays a role in wound healing, tissue remodeling, cartilage degradation, bone development, bone mineralization and ossification. Required for normal embryonic bone development and ossification. Plays a role in the healing of bone fractures via endochondral ossification. Plays a role in wound healing, probably by a mechanism that involves proteolytic activation of TGFB1 and degradation of CTGF. Plays a role in keratinocyte migration during wound healing. May play a role in cell migration and in tumor cell invasion.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505209
NCBI GeneID	4322
NCBI Accession #	NP_002418.1
NCBI GenBank Nucleotide #	NM_002427.3
UniProt Accession #	P45452
Molecular Weight	53,820 Da
NCBI Official Full Name	collagenase 3 preproprotein
NCBI Official Synonym Full Names	matrix metalloproteinase 13 (collagenase 3)
NCBI Official Symbol	MMP13
NCBI Official Synonym Symbols	CLG3; MANDP1
NCBI Protein Information	collagenase 3; MMP-13; matrix metalloproteinase 13 (collagenase 3)
UniProt Protein Name	Collagenase 3
UniProt Gene Name	MMP13
UniProt Synonym Gene Names	MMP-13
UniProt Entry Name	MMP13_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

