

Rabbit anti-Human ATM Polyclonal Antibody | anti-ATM antibody

ATM Antibody (N-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA290715	Rabbit	Human	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ATM; Polyclonal Antibody; ATM Antibody (N-term); Serine-protein kinase ATM; Ataxia telangiectasia mutated; A-T mutated; anti-ATM antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Ig
Specificity	This ATM antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 5~34 amino acids from the N-terminal region of human ATM.
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Sequence Length	3056
Applicable Applications for anti-ATM antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Cellular Location	Nucleus. Cytoplasmic vesicle Note=Primarily nuclear. Found also in endocytic vesicles in association with beta-adaptin.
Tissue Location	Found in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain, heart, spleen, thymus, testis, ovary, small intestine, colon and leukocytes
Antigen Region	5-34
Preparation and Storage	Maintain refrigerated at 2-8°C. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



WB (Western Blot)
Anti-ATM Antibody (C12) at 1:500 dilution + HeLa whole cell lysateLysates/proteins at 20 ug per lane.SecondaryGoat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.Predicted band size : 351 kDaBlocking/Dilution buffer: 5% NFD/MTBST.

ADDITIONAL INFORMATION

Related Product Information for anti-ATM antibody	ATM is involved in signal transduction, cell cycle control and DNA repair, and may function as a tumor suppressor. It is necessary for activation of ABL1 and SAPK, and phosphorylates p53, NFKBIA, BRCA1, CTIP, NIBRIN (NBS1), TERF1, and RAD9. This protein has potential roles in vesicle and/or protein transport, T-cell development, gonad and neurological function. ATM is also part of the BRCA1-associated genome surveillance complex. ATM is induced by ionizing radiation. Defects in ATM are the cause of ataxia talangiectasia (AT), also known as Louis-Bar syndrome, a rare recessive disorder characterized by progressive cerebellar ataxia, dilation of the blood vessels in the conjunctiva and eyeballs, immunodeficiency, growth retardation and sexual immaturity. About 30% of AT patients develop lymphomas and leukemias. Defects in ATM also contribute to T-cell acute lymphoblastic leukemia (TALL) and T-prolymphocytic leukemia (TPLL). TPLL is characterized by a high white blood cell count, with a predominance of prolymphocytes, marked splenomegaly, lymphadenopathy, skin lesions and serous effusion. Defects in ATM also contribute to B-cell non-Hodgkin's lymphomas, and to B-cell chronic lymphocytic leukemia, a disease characterized by accumulation of mature CD5+ B lymphocytes, lymphadenopathy, immunodeficiency and bone marrow failure.
Product Categories/Family for anti-ATM antibody	Primary Antibodies; Cancer; Neuroscience
References	Suzuki, A., et al., J. Biol. Chem. 278 (1):48-53 (2003). Kishi, S., et al., J. Biol. Chem. 276 (31):29282-29291 (2001). Schaffner, C., et al., Proc. Natl. Acad. Sci. U.S.A. 97 (6):2773-2778 (2000). Gatei, M., et al., Nat. Genet. 25 (1):115-119 (2000). Becker-Catania, S.G., et al., Mol. Genet. Metab. 70 (2):122-133 (2000).

NCBI AND UNIPROT INFORMATION

NCBI GI #	71902540
NCBI GeneID	472
NCBI Accession #	NP_000042.3
NCBI GenBank Nucleotide #	NM_000051.3
UniProt Accession #	Q13315; O15429; Q12758; Q16551; Q93007; Q9NP02; Q9UCX7; B2RXN5
NCBI Official Full Name	serine-protein kinase ATM
NCBI Official Synonym Full Names	ATM serine/threonine kinase
NCBI Official Symbol	ATM
NCBI Official Synonym Symbols	AT1; ATA; ATC; ATD; ATE; ATDC; TEL1; TELO1

NCBI Protein Information	serine-protein kinase ATM
UniProt Protein Name	Serine-protein kinase ATM
UniProt Gene Name	ATM
UniProt Synonym Gene Names	A-T mutated
UniProt Entry Name	ATM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/atm-antibody-immunohistochemistry-western-blot-elisa/290715>