

Rabbit NY-CO-9 Polyclonal Antibody | anti-NY-CO-9 antibody

NY-CO-9 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29084	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

NY-CO-9; Polyclonal Antibody; NY-CO-9 Polyclonal Antibody; HDAC5; KIAA0600; Histone deacetylase 5; HD5; Antigen NY-CO-9; anti-NY-CO-9 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-NY-CO-9 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.



Application Data

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.


IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-NY-CO-9 antibody	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation by repressing transcription of myocyte enhancer MEF2C. During muscle differentiation, it shuttles into the cytoplasm, allowing the expression of myocyte enhancer factors. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	62750349
NCBI GeneID	10014
NCBI Accession #	NP_001015053.1
NCBI GenBank Nucleotide #	NM_001015053.1
UniProt Accession #	Q9UQL6; O60340; O60528; Q96DY4; C9JFV9
Molecular Weight	121,978 Da
NCBI Official Full Name	histone deacetylase 5 isoform 3
NCBI Official Synonym Full Names	histone deacetylase 5
NCBI Official Symbol	HDAC5
NCBI Official Synonym Symbols	HD5; NY-CO-9
NCBI Protein Information	histone deacetylase 5; antigen NY-CO-9
UniProt Protein Name	Histone deacetylase 5
UniProt Gene Name	HDAC5
UniProt Synonym Gene Names	KIAA0600; HD5
UniProt Entry Name	HDAC5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.