

Rabbit anti-Human, Mouse STAT3 Polyclonal Antibody | anti-STAT3 antibody

STAT3 (S727) Antibody

CATALOG # AAA290931	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	-----------------------------------	--


PRODUCT SYNONYMS

STAT3; Polyclonal Antibody; STAT3 (S727) Antibody; Signal transducer and activator of transcription 3; Acute-phase response factor; APRF; anti-STAT3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	Ig
Specificity	This STAT3(S727) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 710-740 amino acids from the human region of human STAT3(S727).
Purity/Purification	Purified through a protein A column, followed by peptide affinity purification.
Form/Format	Supplied in PBS with 0.09% (W/V) sodium azide.
Sequence Length	769
Applicable Applications for anti-STAT3 antibody	ELISA, WB (Western Blot)
Antigen Type	Synthesized Peptide
Antigen Source	Human
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: WB - 1:2000Anti-STAT3(S727) Antibody at 1:2000 dilution + NIH/3T3 whole cell lysate Lysates/proteins at 20 ug per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 88 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



WB (Western Blot)

Dilution: WB - 1:500Anti-STAT3(S727) Antibody at 1:500 dilution + 293 whole cell lysate Lysates/proteins at 20 ug per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 88 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-STAT3 antibody	Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF and other growth factors. May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4. Binds to the interleukin-6 (IL-6)- responsive elements identified in the promoters of various acute- phase protein genes. Activated by IL31 through IL31RA. Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity. Plays an important role in host defense in methicillin-resistant S.aureus lung infection by regulating the expression of the antimicrobial lectin REG3G (By similarity).
Product Categories/Family for anti-STAT3 antibody	Stem Cells
References	Akira S.,et al.Cell 77:63-71(1994). Della Pietra L.,et al.Gene 213:119-124(1998). Feinstein E.,et al.Patent number EP2440214, 18-APR-2012. Ota T.,et al.Nat. Genet. 36:40-45(2004). Zody M.C.,et al.Nature 440:1045-1049(2006).

NCBI AND UNIPROT INFORMATION

NCBI GI #	21618338
NCBI GeneID	6774
NCBI Accession #	NP_003141.2
NCBI GenBank Nucleotide #	NM_003150.3
UniProt Accession #	P40763
Molecular Weight	88068 Da
NCBI Official Full Name	signal transducer and activator of transcription 3 isoform 2
NCBI Official Synonym Full Names	signal transducer and activator of transcription 3
NCBI Official Symbol	STAT3
NCBI Official Synonym Symbols	APRF; HIES; ADMIO; ADMIO1
NCBI Protein Information	signal transducer and activator of transcription 3
UniProt Protein Name	Signal transducer and activator of transcription 3
UniProt Gene Name	STAT3
UniProt Synonym Gene Names	APRF
UniProt Entry Name	STAT3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/stat3-antibody-western-blot-elisa/290931>