

Rabbit C/EBP delta/epsilon Polyclonal Antibody | anti-C/EBPdelta/epsilon antibody

Anti-C/EBP delta/epsilon Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291039	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

C/EBP delta/epsilon; Polyclonal Antibody; Anti-C/EBP delta/epsilon Antibody; CEBPD; CCAAT/enhancer-binding protein delta; C/EBP delta; Nuclear factor NF-IL6-beta; NF-IL6-beta; CEBPE; CCAAT/enhancer-binding protein epsilon; C/EBP epsilon; anti-C/EBPdelta/epsilon antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-C/EBPdelta/epsilon antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES


IF (Immunofluorescence)

Dilution: IF: 1/50 - 1/200 Immunofluorescent analysis of C/EBP delta/epsilon staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.


IHC (Immunohistochemistry)

Dilution: IH 1/50 - 1/200 Immunohistochemical analysis of C/EBP delta/epsilon staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Dilution: WB 1/500 - 1/1000 Western blot analysis of C/EBP delta/epsilon expression in CT26 (A), Hela (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-C/EBPdelta/epsilon antibody	Rabbit polyclonal antibody to C/EBP delta/epsilon
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	160332350
NCBI GeneID	1052
UniProt Accession #	P49716; Q14937; Q2M2X9
Molecular Weight	269
NCBI Official Full Name	CCAAT/enhancer-binding protein delta
NCBI Official Synonym Full Names	CCAAT/enhancer binding protein (C/EBP), delta
NCBI Official Symbol	CEBPD
NCBI Official Synonym Symbols	CELF; CRP3; C/EBP-delta; NF-IL6-beta
NCBI Protein Information	CCAAT/enhancer-binding protein delta; c/EBP delta; nuclear factor NF-IL6-beta
UniProt Protein Name	CCAAT/enhancer-binding protein delta
UniProt Gene Name	CEBPD
UniProt Synonym Gene Names	C/EBP delta; NF-IL6-beta
UniProt Entry Name	CEBPD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/c-ebp-delta-epsilon-antibody-immunohistochemistry-immunofluorescence-western-blot-elisa/291039>