

Rabbit EGR Polyclonal Antibody | anti-EGR antibody

Anti-EGR Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291042	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

EGR; Polyclonal Antibody; Anti-EGR Antibody; EGR1; KROX24; ZNF225; Early growth response protein 1; EGR-1; AT225; Nerve growth factor-induced protein A; NGFI-A; Transcription factor ETR103; Transcription factor Zif268; Zinc finger protein 225; Zinc finger protein Krox-24; EGR2; KROX20; E3 SUMO-protein ligase EGR2; AT591; Early growth response protein 2; EGR-2; Zinc finger protein Krox-20; anti-EGR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-EGR antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IF (Immunofluorescence)

Dilution: IF: 1/50 - 1/200 Immunofluorescent analysis of EGR staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Dilution: IH 1/50 - 1/200 Immunohistochemical analysis of EGR staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Dilution: WB 1/500 - 1/1000 Western blot analysis of EGR expression in Hela (A), PC3 (B), rat ovary (C), MCF7 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-EGR antibody	Rabbit polyclonal antibody to EGR
---	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503493
-----------	---------

NCBI GeneID	1958
NCBI Accession #	NP_001955.1
NCBI GenBank Nucleotide #	NM_001964.2
UniProt Accession #	P18146
Molecular Weight	57,507 Da
NCBI Official Full Name	early growth response protein 1
NCBI Official Synonym Full Names	early growth response 1
NCBI Official Symbol	EGR1
NCBI Official Synonym Symbols	TIS8; AT225; G0S30; NGFI-A; ZNF225; KROX-24; ZIF-268
NCBI Protein Information	early growth response protein 1; EGR-1; zinc finger protein 225; transcription factor ETR103; transcription factor Zif268; zinc finger protein Krox-24; nerve growth factor-induced protein A
UniProt Protein Name	Early growth response protein 1
UniProt Gene Name	EGR1
UniProt Synonym Gene Names	KROX24; ZNF225; EGR-1; NGFI-A
UniProt Entry Name	EGR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/egr-antibody-immunohistochemistry-immunofluorescence-western-blot-elisa/291042>