

Rabbit Rho A Polyclonal Antibody | anti-RHOA antibody

Rho A Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29110	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

Rho A; Polyclonal Antibody; Rho A Polyclonal Antibody; RHOA; ARH12; ARHA; RHO12; Transforming protein RhoA; Rho cDNA clone 12; h12; anti-RHOA antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-RHOA antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES


IHC (Immunohistochemistry)


Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.


IHC (Immunohistochemistry)



Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.


IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.


IF (Immunofluorescence)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.



Application Data

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Application Data

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-RHOA antibody

Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers. Involved in a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Plays an essential role in cleavage furrow formation. Required for the apical junction formation of keratinocyte cell-cell adhesion. Stimulates PKN2 kinase activity. May be an activator of PLCE1. Activated by ARHGEF2, which promotes the exchange of GDP for GTP. Essential for the SPATA13-mediated regulation of cell migration and adhesion assembly and disassembly. The MEMO1-RHOA-DIAPH1 signaling pathway plays an important role in ERBB2-dependent stabilization of microtubules at the cell cortex. It controls the localization of APC and CLASP2 to the cell membrane, via the regulation of GSK3B activity. In turn, membrane-bound APC allows the localization of the MACF1 to the cell membrane, which is required for microtubule capture and stabilization. Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers. Involved in a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Plays an essential role in cleavage furrow formation. Required for the apical junction formation of keratinocyte cell-cell adhesion. May be an activator of PLCE1. Activated by ARHGEF2, which promotes the exchange of GDP for GTP. Essential for the SPATA13-mediated regulation of cell migration and adhesion assembly and disassembly. The MEMO1-RHOA-DIAPH1 signaling pathway plays an important role in ERBB2-dependent stabilization of microtubules at the cell cortex. It controls the localization of APC and CLASP2 to the cell membrane, via the regulation of GSK3B activity. In turn, membrane-bound APC allows the localization of the MACF1 to the cell membrane, which is required for microtubule capture and stabilization (By similarity). Regulates KCNA2 potassium channel activity by reducing its location at the cell surface in response to CHRM1 activation; promotes KCNA2 endocytosis (PubMed:19403695).

NCBI AND UNIPROT INFORMATION

NCBI GI #	47606458
NCBI GeneID	387
UniProt Accession #	P61586 ; P06749 ; Q53HM4 ; Q5U024 ; Q9UDJ0 ; Q9UEJ4
Molecular Weight	193
NCBI Official Full Name	Transforming protein RhoA
NCBI Official Synonym Full Names	ras homolog family member A
NCBI Official Symbol	RHOA
NCBI Official Synonym Symbols	ARHA; ARH12; RHO12; RHOH12
NCBI Protein Information	transforming protein RhoA; h12; oncogene RHO H12; rho cDNA clone 12; Aplysia ras-related homolog 12; small GTP binding protein RhoA; ras homolog gene family, member A
UniProt Protein Name	Transforming protein RhoA
UniProt Gene Name	RHOA
UniProt Synonym Gene Names	ARH12; ARHA; RHO12; h12
UniProt Entry Name	RHOA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

