

Rabbit PAK1/2/3 Polyclonal Antibody | anti-PAK1/2/3 antibody

Anti-PAK1/2/3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291125	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

PAK1/2/3; Polyclonal Antibody; Anti-PAK1/2/3 Antibody; PAK1; Serine/threonine-protein kinase PAK 1; Alpha-PAK; p21-activated kinase 1; PAK-1; p65-PAK; PAK2; Serine/threonine-protein kinase PAK 2; Gamma-PAK; PAK65; S6/H4 kinase; p21-activated kinase 2; PAK-2; p58; PAK3; OPHN3; Serine/threonine-protein kinase PAK 3; Beta-PAK; Oligophrenin-3; p21-activated kinase 3; PAK-3; anti-PAK1/2/3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-PAK1/2/3 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IF (Immunofluorescence)

Dilution: IF: 1/50 - 1/200 Immunofluorescent analysis of PAK1/2/3 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Dilution: IH 1/50 - 1/200 Immunohistochemical analysis of PAK1/2/3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Dilution: WB 1/500 - 1/1000 Western blot analysis of PAK1/2/3 expression in A375 (A), A549 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PAK1/2/3 antibody	Rabbit polyclonal antibody to PAK1/2/3
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	190886457
NCBI GeneID	5058
NCBI Accession #	NP_001122092.1
NCBI GenBank Nucleotide #	NM_001128620.1
UniProt Accession #	Q13153; O75561; Q13567; Q32M53; Q32M54; Q86W79
Molecular Weight	60,647 Da
NCBI Official Full Name	serine/threonine-protein kinase PAK 1 isoform 1
NCBI Official Synonym Full Names	p21 protein (Cdc42/Rac)-activated kinase 1
NCBI Official Symbol	PAK1
NCBI Official Synonym Symbols	PAKalpha
NCBI Protein Information	serine/threonine-protein kinase PAK 1; p65-PAK; alpha-PAK; STE20 homolog, yeast; p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related); p21/Cdc42/Rac1-activated kinase 1 (STE20 homolog, yeast)
UniProt Protein Name	Serine/threonine-protein kinase PAK 1
UniProt Gene Name	PAK1
UniProt Synonym Gene Names	PAK-1
UniProt Entry Name	PAK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pak1-2-3-antibody-immunohistochemistry-immunofluorescence-western-blot-elisa/291125>