

Rabbit EPHB1/2 Polyclonal Antibody | anti-EPHB1/2 antibody

Anti-EPHB1/2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291250	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

EPHB1/2; Polyclonal Antibody; Anti-EPHB1/2 Antibody; EPHB1; ELK; EPHT2; HEK6; NET; Ephrin type-B receptor 1; EPH tyrosine kinase 2; EPH-like kinase 6; EK6; hEK6; Neuronally-expressed EPH-related tyrosine kinase; Tyrosine-protein kinase receptor EPH-2; EPHB2; DRT; EPHT3; EPTH3; ERK; HEK5; TYRO5; Ephrin type-B receptor 2; Developmentally-regulated Eph-related tyrosine kinase; ELK-related tyrosine kinase; EPH tyrosine kinase 3; EPH-like kinase 5; EK5; hEK5; Renal carcinoma antigen NY-REN-47; Tyrosine-protein kinase TYRO5; Tyrosine-protein kinase receptor EPH-3; anti-EPHB1/2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-EPHB1/2 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IF (Immunofluorescence)

Dilution: IF: 1/50 - 1/200 Immunofluorescent analysis of EPHB1/2 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Dilution: IH 1/50 - 1/200 Immunohistochemical analysis of EPHB1/2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Dilution: WB 1/500 - 1/1000 Western blot analysis of EPHB1/2 expression in U87MG (A), rat brain (B), mouse heart (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-EPHB1/2 antibody

Rabbit polyclonal antibody to EPHB1/2

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758284
NCBI GeneID	2047
NCBI Accession #	NP_004432.1
NCBI GenBank Nucleotide #	NM_004441.4
UniProt Accession #	P54762; O43569; O95142; O95143; Q0VG87; A8K593
Molecular Weight	109,885 Da
NCBI Official Full Name	ephrin type-B receptor 1
NCBI Official Synonym Full Names	EPH receptor B1
NCBI Official Symbol	EPHB1
NCBI Official Synonym Symbols	ELK; NET; Hek6; EPHT2
NCBI Protein Information	ephrin type-B receptor 1; EK6; EPH-like kinase 6; eph tyrosine kinase 2; soluble EPHB1 variant 1; tyrosine-protein kinase receptor EPH-2; neuronally-expressed EPH-related tyrosine kinase
UniProt Protein Name	Ephrin type-B receptor 1
UniProt Gene Name	EPHB1
UniProt Synonym Gene Names	ELK; EPHT2; HEK6; NET; EK6; hEK6; NET
UniProt Entry Name	EPHB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ephb1-2-antibody-immunohistochemistry-immunofluorescence-western-blot-elisa/291250>