

Rabbit HOXc8 Polyclonal Antibody | anti-HOXc8 antibody

HOXc8 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291265	Rabbit	Rat, Pig, Cow	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HOXc8; Polyclonal Antibody; HOXc8 Polyclonal Antibody; Homeobox protein Hox-C8; Homeobox protein Hox-3A; HOXC8; HOX3A; anti-HOXc8 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-HOXc8 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Human cervical cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Histone H3) Polyclonal Antibody, Unconjugated (bs-0394R) at 1:500 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Human cervical cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Hox-3.1) Polyclonal Antibody, Unconjugated (bs-0394R) at 1:500 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded Rat small intestine; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer normal goat serum at 37 degree C for 30min; Antibody incubation with (Hox-3.1) Polyclonal Antibody, Unconjugated (bs-0394R) at 1:500 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



WB (Western Blot)

Sample:Brain(Rat) lysate at 30ug;Liver(Rat) lysate at 30ug;Primary: Anti-HOXc8 (bs-0394R) at 1:200;Secondary: HRP conjugated Goat-Anti-Rabbit IgG(bs-0295G) at 1: 3000;Predicted band size : 27kD Observed band size : 27kD, 43kD

NCBI AND UNIPROT INFORMATION

NCBI GI #	13124745
NCBI GeneID	3224

UniProt Accession #	P31273; O15221; O15362; A8K4J4
Molecular Weight	242
NCBI Official Full Name	Homeobox protein Hox-C8
NCBI Official Synonym Full Names	homeobox C8
NCBI Official Symbol	HOXC8
NCBI Official Synonym Symbols	HOX3; HOX3A
NCBI Protein Information	homeobox protein Hox-C8; homeo box 3A; homeo box C8; homeobox protein Hox-3A; Hox-3.1, mouse, homolog of
UniProt Protein Name	Homeobox protein Hox-C8
UniProt Gene Name	HOXC8
UniProt Synonym Gene Names	HOX3A
UniProt Entry Name	HXC8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hoxc8-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291265>